

The Project Gutenberg eBook of Elliptic Functions, by Arthur L. Baker

This eBook is for the use of anyone anywhere at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at [www.gutenberg.org](http://www.gutenberg.org)

Title: Elliptic Functions  
An Elementary Text-Book for Students of Mathematics

Author: Arthur L. Baker

Release Date: January 25, 2010 [EBook #31076]

Language: English

Character set encoding: ISO-8859-1

\*\*\* START OF THIS PROJECT GUTENBERG EBOOK ELLIPTIC FUNCTIONS \*\*\*

Produced by Andrew D. Hwang, Brenda Lewis and the Online Distributed Proofreading Team at <http://www.pgdp.net> (This file was produced from images from the Cornell University Library: Historical Mathematics Monographs collection.)

#### TRANSCRIBER'S NOTE

This book was produced from images provided by the Cornell University Library: Historical Mathematics Monographs collection.

Minor typographical corrections and presentational changes have been made without comment. The calculations preceding equation (15) on [page 12](#) (page 12 of the original) have been re-formatted.

This PDF file is optimized for screen viewing, but may easily be recompiled for printing. Please see the preamble of the  $\text{\LaTeX}$  source file for instructions.

# ELLIPTIC FUNCTIONS.

AN ELEMENTARY TEXT-BOOK FOR  
STUDENTS OF MATHEMATICS.

BY

ARTHUR L. BAKER, C.E., PH.D.,

PROFESSOR OF MATHEMATICS IN THE STEVENS SCHOOL OF THE STEVENS INSTITUTE OF  
TECHNOLOGY, HOBOKEN, N. J.; FORMERLY PROFESSOR IN THE PARDEE  
SCIENTIFIC DEPARTMENT, LAFAYETTE COLLEGE, EASTON, PA.

---

$$\sin \operatorname{am} u = \frac{1}{\sqrt{k}} \cdot \frac{H(u)}{\Theta(u)}.$$

---

NEW YORK:  
JOHN WILEY & SONS,  
53 EAST TENTH STREET.  
1890.

Copyright, 1890,  
BY  
Arthur L. Baker.

ROBERT DRUMMOND,  
*Electrotyper,*  
444 & 446 Pearl Street,  
New York.

FERRIS BROS.,  
*Printers,*  
326 Pearl Street,  
New York.

# PREFACE.

---

IN the works of Abel, Euler, Jacobi, Legendre, and others, the student of Mathematics has a most abundant supply of material for the study of the subject of Elliptic Functions.

These works, however, are not accessible to the general student, and, in addition to being very technical in their treatment of the subject, are moreover in a foreign language.

It is in the hope of smoothing the road to this interesting and increasingly important branch of Mathematics, and of putting within reach of the English student a tolerably complete outline of the subject, clothed in simple mathematical language and methods, that the present work has been compiled.

New or original methods of treatment are not to be looked for. The most that can be expected will be the simplifying of methods and the reduction of them to such as will be intelligible to the average student of Higher Mathematics.

I have endeavored throughout to use only such methods as are familiar to the ordinary student of Calculus, avoiding those methods of discussion dependent upon the properties of double periodicity, and also those depending upon Functions of Complex Variables. For the same reason I have not carried the discussion of the  $\Theta$  and  $H$  functions further.

Among the minor helps to simplicity is the use of zero subscripts to indicate decreasing series in the Landen Transformation, and of numerical subscripts to indicate increasing series. I have adopted the notation of Gudermann, as being more simple than that of Jacobi.

I have made free use of the following works: JACOBI'S *Fundamenta Nova Theoriæ Func. Ellip.*; HOUËL'S *Calcul Infinitésimal*;

LEGENDRE'S *Traité des Fonctions Elliptiques*; DUREGE'S *Theorie der Elliptischen Functionen*; HERMITE'S *Théorie des Fonctions Elliptiques*; VERHULST'S *Théorie des Fonctions Elliptiques*; BERTRAND'S *Calcul Intégral*; LAURENT'S *Théorie des Fonctions Elliptiques*; CAYLEY'S *Elliptic Functions*; BYERLY'S *Integral Calculus*; SCHLOMILCH'S *Die Höheren Analysis*; BRIOT ET BOUQUET'S *Fonctions Elliptiques*.

I have refrained from any reference to the Gudermann or Weierstrass functions as not within the scope of this work, though the Gudermannians might have been interesting examples of verification formulæ. The arithmetico-geometrical mean, the march of the functions, and other interesting investigations have been left out for want of room.

# CONTENTS

---

INTRODUCTORY CHAPTER. . . . .	PAGE
CHAP. I. ELLIPTIC INTEGRALS. . . . .	4
II. ELLIPTIC FUNCTIONS. . . . .	16
III. PERIODICITY OF THE FUNCTIONS. . . . .	24
IV. LANDEN'S TRANSFORMATION . . . . .	33
V. COMPLETE FUNCTIONS . . . . .	50
VI. EVALUATION FOR $\phi$ . . . . .	53
VII. DEVELOPMENT OF ELLIPTIC FUNCTIONS INTO FACTORS. . . . .	56
VIII. THE $\Theta$ FUNCTION. . . . .	71
IX. THE $\Theta$ AND $H$ FUNCTIONS. . . . .	74
X. ELLIPTIC INTEGRALS OF THE SECOND ORDER. . . . .	86
XI. ELLIPTIC INTEGRALS OF THE THIRD ORDER. . . . .	96
XII. NUMERICAL CALCULATIONS. $q$ . . . . .	101
XIII. NUMERICAL CALCULATIONS. $K$ . . . . .	105
XIV. NUMERICAL CALCULATIONS. $u$ . . . . .	111
XV. NUMERICAL CALCULATIONS. $\phi$ . . . . .	119
XVI. NUMERICAL CALCULATIONS. $E(k, \phi)$ . . . . .	123
XVII. APPLICATIONS. . . . .	128

# ELLIPTIC FUNCTIONS.

---

## INTRODUCTORY CHAPTER.\*

THE first step taken in the theory of Elliptic Functions was the determination of a relation between the amplitudes of three functions of either order, such that there should exist an algebraic relation between the three functions themselves of which these were the amplitudes. It is one of the most remarkable discoveries which science owes to Euler. In 1761 he gave to the world the complete integration of an equation of two terms, each an elliptic function of the first or second order, not separately integrable.

This integration introduced an arbitrary constant in the form of a third function, related to the first two by a given equation between the amplitudes of the three.

In 1775 Landen, an English mathematician, published his celebrated theorem showing that any arc of a hyperbola may be measured by two arcs of an ellipse, an important element of the theory of Elliptic Functions, but *then* an isolated result. The great problem of comparison of Elliptic Functions of different moduli remained unsolved, though Euler, in a measure, exhausted the comparison of functions of the same modulus. It was completed in 1784 by Lagrange, and for the computation of numerical results leaves little to be desired. The value of a function may be determined by it, in terms of increasing or diminishing moduli,

---

\*Condensed from an article by Rev. Henry Moseley, M.A., F.R.S., Prof. of Nat. Phil. and Ast., King's College, London.



until at length it depends upon a function having a modulus of zero, or unity.

For all practical purposes this was sufficient. The enormous task of calculating tables was undertaken by Legendre. His labors did not end here, however. There is none of the discoveries of his predecessors which has not received some perfection at his hands; and it was he who first supplied to the whole that connection and arrangement which have made it an independent science.

The theory of Elliptic Integrals remained at a standstill from 1786, the year when Legendre took it up, until the year 1827, when the second volume of his *Traité des Fonctions Elliptiques* appeared. Scarcely so, however, when there appeared the researches of Jacobi, a Professor of Mathematics in Königsberg, in the 123d number of the *Journal of Schumacher*, and those of Abel, Professor of Mathematics at Christiania, in the 3d number of *Crelle's Journal* for 1827.

These publications put the theory of Elliptic Functions upon an entirely new basis. The researches of Jacobi have for their principal object the development of that general relation of functions of the first order having different moduli, of which the scales of Lagrange and Legendre are particular cases.

It was to Abel that the idea first occurred of treating the Elliptic Integral as a function of its amplitude. Proceeding from this new point of view, he embraced in his speculations all the principal results of Jacobi. Having undertaken to develop the principle upon which rests the fundamental proposition of Euler establishing an algebraic relation between three functions which have the same moduli, dependent upon a certain relation of their amplitudes, he has extended it from three to an indefinite number of functions; and from Elliptic Functions to an infinite number of other functions embraced under an indefinite number of classes, of which that of Elliptic Functions is but one; and each class having a division analogous to that of Elliptic Functions into three

orders having common properties.

The discovery of Abel is of infinite moment as presenting the first step of approach towards a more complete theory of the infinite class of ultra elliptic functions, destined probably ere long to constitute one of the most important of the branches of transcendental analysis, and to include among the integrals of which it effects the solution some of those which at present arrest the researches of the philosopher in the very elements of physics.

# CHAPTER I.

## ELLIPTIC INTEGRALS.

THE integration of irrational expressions of the form

$$X dx \sqrt{A + Bx + Cx^2},$$

or

$$\frac{X dx}{\sqrt{A + Bx + Cx^2}},$$

$X$  being a rational function of  $x$ , is fully illustrated in most elementary works on Integral Calculus, and shown to depend upon the transcendentials known as logarithms and circular functions, which can be calculated by the proper logarithmic and trigonometric tables.

When, however, we undertake to integrate irrational expressions containing higher powers of  $x$  than the square, we meet with insurmountable difficulties. This arises from the fact that the integral sought depends upon a new set of transcendentials, to which has been given the name of *elliptic functions*, and whose characteristics we will learn hereafter.

The name of Elliptic Integrals has been given to the simple integral forms to which can be reduced all integrals of the form

$$(1) \quad V = \int F(X, R) dx,$$

where  $F(X, R)$  designates a rational function of  $x$  and  $R$ , and  $R$  represents a radical of the form

$$R = \sqrt{Ax^4 + Bx^3 + Cx^2 + Dx + E},$$

where  $A, B, C, D, E$  indicate constant coefficients.

We will show presently that all cases of Eq. (1) can be reduced to the three typical forms

$$(2) \quad \begin{aligned} & \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \\ & \int_0^x \frac{x^2 dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \\ & \int_0^x \frac{dx}{(x^2+a)\sqrt{(1-x^2)(1-k^2x^2)}}, \end{aligned}$$

which are called elliptic integrals of the first, second, and third order.

Why they are called *Elliptic* Integrals we will learn further on. The transcendental functions which depend upon these integrals, and which will be discussed in [Chapter IV](#), are called *Elliptic Functions*.

The most general form of Eq. (1) is

$$(3) \quad V = \int \frac{A + BR}{C + DR} dx;$$

where  $A, B, C,$  and  $D$  stand for rational integral functions of  $x$ .

$\frac{A + BR}{C + DR}$  can be written

$$\begin{aligned} \frac{A + BR}{C + DR} &= \frac{AC - BDR^2}{C^2 - D^2R^2} - \frac{(AD - CB)R^2}{C^2 - D^2R^2} \cdot \frac{1}{R} \\ &= N - \frac{P}{R}; \end{aligned}$$

$N$  and  $P$  being rational integral functions of  $x$ . Whence Eq. (3) becomes

$$(4) \quad V = \int N dx - \int \frac{P dx}{R}.$$

Eq. (4) shows that the most general form of  $V$  can be made to depend upon the expressions

$$(5) \quad V' = \int \frac{P dx}{R},$$

and

$$\int N dx.$$

This last form is rational, and needs no discussion here. We can write

$$\begin{aligned} P &= \frac{G_0 + G_1x + G_2x^2 + \cdots}{H_0 + H_1x + H_2x^2 + \cdots} \\ &= \frac{G_0 + G_2x^2 + G_4x^4 + \cdots + (G_1 + G_3x^2 + \cdots)x}{H_0 + H_2x^2 + H_4x^4 + \cdots + (H_1 + H_3x^2 + \cdots)x}. \end{aligned}$$

Multiplying both numerator and denominator by

$$H_0 + H_2x^2 + H_4x^4 + \cdots - (H_1 + H_3x^2 + H_5x^4 + \cdots)x,$$

we have a new denominator which contains only powers of  $x^2$ . The result takes the following form:

$$\begin{aligned} P &= \frac{M_0 + M_2x^2 + M_4x^4 + \cdots + (M_1 + M_3x^2 + M_5x^4 + \cdots)x}{N_0 + N_2x^2 + N_4x^4 + N_6x^6 + \cdots} \\ &= \Phi(x^2) + \Psi(x^2) \cdot x. \end{aligned}$$

Equation (5) thus becomes

$$(6) \quad V' = \int \frac{\Phi(x^2) dx}{R} + \int \frac{\Psi(x^2) \cdot x \cdot dx}{R}.$$

We shall see presently that  $R$  can always be assumed to be of the form

$$\sqrt{(1-x^2)(1-k^2x^2)}.$$

Therefore, putting  $x^2 = z$ , the second integral in Eq. (6) takes the form

$$\frac{1}{2} \int \frac{\Psi(z) \cdot dz}{\sqrt{(1-z)(1-k^2z)}},$$

which can be integrated by the well-known methods of Integral Calculus, resulting in logarithmic and circular transcendentals.

There remains, therefore, only the form

$$\int \frac{\Phi(x^2) dx}{R}$$

to be determined.

We will now show that  $R$  can always be assumed to be in the form

$$\sqrt{(1-x^2)(1-k^2x^2)}.$$

We have

$$\begin{aligned} R &= \sqrt{Ax^4 + Bx^3 + Cx^2 + Dx + E} \\ &= \sqrt{G(x-a)(x-b)(x-c)(x-d)}, \end{aligned}$$

$a, b, c$ , and  $d$  being the roots of the polynomial of the fourth degree, and  $G$  any number, real or imaginary, depending upon the coefficients in the given polynomial.

Substituting in [equation \(1\)](#)

$$x = \frac{p + qy}{1 + y},$$

we have

$$(7) \quad V = \int \phi(y, \rho) dy,$$

$\rho$  designating the radical

$$\rho = \sqrt{G[p - a + (q - a)y][p - b + (q - b)y][p - c + (q - c)y] \cdots}$$

In order that the odd powers of  $y$  under the radical may disappear we must have their coefficients equal to zero; i.e.,

$$\begin{aligned} (p - a)(q - b) + (p - b)(q - a) &= 0, \\ (p - c)(q - d) + (p - d)(q - c) &= 0; \end{aligned}$$

whence

$$\begin{aligned} 2pq - (p + q)(a + b) + 2ab &= 0, \\ 2pq - (p + q)(c + d) + 2cd &= 0, \end{aligned}$$

and

$$(8) \quad \begin{cases} pq = \frac{ab(c + d) - cd(a + b)}{a + b - (c + d)}, \\ p + q = \frac{2ab - 2cd}{a + b - (c + d)}. \end{cases}$$

Equation (8) shows that  $p$  and  $q$  are real quantities, whether the roots  $a$ ,  $b$ ,  $c$ , and  $d$  are real or imaginary;  $a$ ,  $b$ , and  $c$ ,  $d$  being the conjugate pairs.

Hence equation (1) can always be reduced to the form of equation (7), which contains only the second and fourth powers of the variable.

This transformation seems to fail when  $a + b - (c + d) = 0$ ; but in that case we have

$$R = \sqrt{G[x^2 - (a + b)x + ab][x^2 - (a + b)x + cd]},$$

and substituting

$$x = y - \frac{a+b}{2}$$

will cause the odd powers of  $y$  to disappear as before.

If the radical should have the form

$$\sqrt{G(x-a)(x-b)(x-c)},$$

placing  $x = y^2 + a$ , we get

$$V = \int \phi(y, \rho) dy,$$

$$\rho = \sqrt{G(y^2 + a - b)(y^2 + a - c)},$$

$\phi$  designating a rational function of  $y$  and  $\rho$ .

Thus all integrals of the form contained in [equation \(1\)](#), in which  $R$  stands for a quadratic surd of the third or fourth degree, can be reduced to the form

$$(9) \quad V = \int \phi(x, R) dx,$$

$R$  designating a radical of the form

$$\sqrt{G(1+mx^2)(1+nx^2)},$$

$m$  and  $n$  designating constants.

It is evident that if we put

$$x' = x\sqrt{-m}, \quad k^2 = -\frac{n}{m},$$

we can reduce the radical to the form

$$\sqrt{(1-x^2)(1-k^2x^2)}.$$



We shall see later on that the quantity  $k^2$ , to which has been given the name *modulus*, can always be considered real and less than unity.

Combining these results with [equation \(6\)](#), we see that the integration of [equation \(1\)](#) depends finally upon the integration of the expression

$$(10) \quad V'' = \int \frac{\phi(x^2) dx}{\sqrt{(1-x^2)(1-k^2x^2)}} = \int \frac{\phi(x^2) dx}{R}.$$

The most general form of  $\phi(x^2)$  is

$$\begin{aligned} \phi(x^2) &= \frac{M_0 + M_2x^2 + M_4x^4 + \dots}{N_0 + N_2x^2 + N_4x^4 + \dots} \\ &= P_0 + P_2x^2 + P_4x^4 + P_6x^6 + \dots \\ &\quad + \sum \frac{L}{(x^2 + a)^n}. \end{aligned}$$

Hence

$$(11) \quad V'' = \sum P \int \frac{x^{2m} dx}{R} + \sum L \int \frac{dx}{(x^2 + a)^n R}.$$

But  $\int \frac{x^{2m} dx}{R}$  depends upon  $\int \frac{dx}{R}$  and  $\int \frac{x^2 dx}{R}$ , which can be shown as follows:

Differentiating  $Rx^{2m-3}$ , we have

$$\begin{aligned} d[x^{2m-3}R] &= d \left[ x^{2m-3} \sqrt{\alpha + \beta x^2 + \gamma x^4} \right] \\ &= (2m-3)x^{2m-4} dx \sqrt{\alpha + \beta x^2 + \gamma x^4} + \frac{x^{2m-3}(\beta x + 2\gamma x^3) dx}{\sqrt{\alpha + \beta x^2 + \gamma x^4}}. \end{aligned}$$

Integrating and collecting, we get

$$\begin{aligned}
 Rx^{2m-3} &= (2m-3)\alpha \int \frac{x^{2m-4} dx}{R} + (2m-2)\beta \int \frac{x^{2m-2} dx}{R} \\
 &\quad + (2m-1)\gamma \int \frac{x^{2m} dx}{R} \\
 (12) \quad &= \alpha' \int \frac{x^{2m-4} dx}{R} + \beta' \int \frac{x^{2m-2} dx}{R} + \gamma' \int \frac{x^{2m} dx}{R}.
 \end{aligned}$$

Whence we get, by taking  $m = 2$ ,

$$(13) \quad Rx = \alpha \int \frac{dx}{R} + \beta \int \frac{x^2 dx}{R} + \gamma \int \frac{x^4 dx}{R},$$

which shows that the general expression  $\int \frac{x^{2m} dx}{R}$  can be found by successive calculations, when we are able to integrate the expressions

$$\int \frac{dx}{R} \quad \text{and} \quad \int \frac{x^2 dx}{R},$$

the first and second of [equation \(2\)](#).

We will now consider the second class of terms in [eq. \(11\)](#), viz.,

$$\frac{L dx}{(x^2 + a)^n R}.$$

This second term is as follows:

$$\begin{aligned}
 (14) \quad \sum \int \frac{L}{(x^2 + a)^n R} &= \int \frac{A dx}{(x^2 + a)^n R} + \int \frac{B dx}{(x^2 + a)^{n-1} R} \\
 &\quad + \int \frac{C dx}{(x^2 + a)^{n-2} R} + \dots
 \end{aligned}$$

Each of these terms can be shown to depend ultimately upon terms of the form

$$\frac{x^2 dx}{R}, \quad \frac{dx}{R}, \quad \text{and} \quad \frac{dx}{(x^2 + a) R}.$$

The two former will be recognized as the two ultimate forms already discussed, the first and second of [equation \(2\)](#). The third is the third one of [equation \(2\)](#).

This dependence of [equation \(14\)](#) can be shown as follows:

We have

$$\begin{aligned} d \left[ \frac{xR}{(x^2+a)^{n-1}} \right] &= \frac{(x^2+a)^{n-1}(x dR + R dx) - 2x^2R(n+1)(x^2+a)^{n-2} dx}{(x^2+a)^{2n-2}} \\ &= \frac{(x^2+a)(x dR + R dx) - 2x^2R(n-1) dx}{(x^2+a)^n}. \end{aligned}$$

Substituting the value of

$$R = \sqrt{\alpha + \beta x^2 + \gamma x^4} \quad \text{and} \quad dR = (\beta x + 2\gamma x^3) \frac{dx}{R},$$

we get

$$\begin{aligned} &d \left[ \frac{xR}{(x^2+a)^{n-1}} \right] \\ &= \frac{(x^2+a)(\beta x^2 + 2\gamma x^4 + \alpha + \beta x^2 + \gamma x^4) - 2x^2(n-1)(\alpha + \beta x^2 + \gamma x^4)}{(x^2+a)^n} \cdot \frac{dx}{R} \\ &= \frac{\left\{ (3\gamma - 2(n-1)\gamma)x^6 + (2\beta + 3a\gamma - 2(n-1)\beta)x^4 \right.}{(x^2+a)^n} \cdot \frac{dx}{R} \\ &\quad \left. + (2a\beta + \alpha - 2(n-1)\alpha)x^2 + a\alpha \right\}}{(x^2+a)^n} \cdot \frac{dx}{R} \\ &= \frac{-(2n-5)\gamma x^6 + (-(2n-4)\beta + 3a\gamma)x^4 + (-(2n-3)\alpha + 2a\beta)x^2 + a\alpha}{(x^2+a)^n} \cdot \frac{dx}{R}; \end{aligned}$$

or, by substituting in the numerator  $x^2 = z - a$ ,

$$= \frac{\left\{ \begin{array}{l} -(2n-5)\gamma z^3 \\ + ((2n-5)3a\gamma - (2n-4)\beta + 3a\gamma)z^2 \\ + (-(2n-5)3a^2\gamma + (2n-4)2a\beta - 6a^2\gamma - (2n-3)\alpha + 2a\beta)z \\ + ((2n-5)a^3\gamma - (2n-4)a^2\beta + 3a^3\gamma + (2n-3)a\alpha - 2a^2\beta + a\alpha) \end{array} \right\}}{(x^2+a)^n} \cdot \frac{dx}{R};$$

or, after resubstituting  $z = x^2 + a$ , and integrating,

$$\begin{aligned} (15) \quad \frac{xR}{(x^2+a)^{n-1}} &= -(2n-5)\gamma \int \frac{dx}{(x^2+a)^{n-3}R} \\ &\quad - (2n-4)(\beta - 3a\gamma) \int \frac{dx}{(x^2+a)^{n-2}R} \\ &\quad - (2n-3)(3a^2\gamma - 2a\beta + \alpha) \int \frac{dx}{(x^2+a)^{n-1}R} \\ &\quad + (2n-2)(a^3\gamma - a^2\beta + a\alpha) \int \frac{dx}{(x^2+a)^n R}. \\ &= \alpha_1 \int \frac{dx}{(x^2+a)^{n-3}R} + \beta_1 \int \frac{dx}{(x^2+a)^{n-2}R} + \gamma_1 \int \frac{dx}{(x^2+a)^{n-1}R} \\ &\quad + \delta_1 \int \frac{dx}{(x^2+a)^n R}. \end{aligned}$$

Making  $n = 2$ , we have

$$\begin{aligned} (16) \quad \frac{xR}{(x^2+a)^1} &= \alpha_1 \int \frac{(x^2+a) dx}{R} + \beta_1 \int \frac{dx}{R} + \gamma_1 \int \frac{dx}{(x^2+a)R} \\ &\quad + \delta_1 \int \frac{dx}{(x^2+a)^2 R}. \end{aligned}$$

Equation (16) shows that

$$\int \frac{dx}{(x^2 + a)^2 R}$$

depends upon the three forms

$$\int \frac{x^2 dx}{R}, \quad \int \frac{dx}{R}, \quad \text{and} \quad \int \frac{dx}{(x^2 + a)R'}$$

the three types of equation (2), and equation (15) shows that the general form

$$\int \frac{dx}{(x^2 + a)^n R}$$

depends ultimately upon the same three types.

We have now discussed every form which the general equation (1) can assume, and shown that they all depend ultimately upon one or more of the three types contained in equation (2).

These three types are called the three Elliptic Integrals of the first, second, and third kind, respectively.

Legendre puts  $x = \sin \phi$ , and reduces the three integrals to the following forms:

$$(17) \quad F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}};$$

$$\frac{1}{k^2} \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} - \frac{1}{k^2} \int_0^\phi \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi;$$

$$(18) \quad \Pi(n, k, \phi) = \int_0^\phi \frac{d\phi}{(1 - n \sin^2 \phi) \sqrt{1 - k^2 \sin^2 \phi}};$$

the first being Legendre's integral of the first kind; the form

$$(19) \quad E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi$$

being the integral of the second kind; and the third one being the integral of the third kind.

The form of the integral of the second kind shows why they are called Elliptic Integrals, the arc of an elliptic quadrant being equal to

$$a \int_0^{\frac{\pi}{2}} \sqrt{1 - e^2 \sin^2 \phi} \cdot d\phi,$$

$\phi$  being the complement of the eccentric angle.

By easy substitutions, we get from Eqs. (17), (18), and (19) the following solutions:

$$\int_0^\phi \frac{\sin^2 \phi}{\Delta} d\phi = \frac{F - E}{k^2};$$

$$\int_0^\phi \frac{\cos^2 \phi}{\Delta} d\phi = \frac{E - (1 - k^2)F}{k^2};$$

$$\int_0^\phi \frac{\tan^2 \phi}{\Delta} d\phi = \frac{\Delta \tan \phi - E}{1 - k^2};$$

$$\int_0^\phi \frac{\sec^2 \phi}{\Delta} d\phi = \frac{\Delta \tan \phi + (1 - k^2)F - E}{1 - k^2};$$

$$\int_0^\phi \frac{1}{\Delta^3} d\phi = \frac{1}{1 - k^2} \left( E - \frac{k^2 \sin \phi \cos \phi}{\Delta} \right);$$

$$\int_0^\phi \frac{\sin^2 \phi}{\Delta^3} d\phi = \frac{1}{1 - k^2} \left( \frac{E - (1 - k^2)F}{k^2} - \frac{\sin \phi \cos \phi}{\Delta} \right);$$

$$\int_0^\phi \frac{\cos^2 \phi}{\Delta^3} d\phi = \frac{F - E}{k^2} + \frac{\sin \phi \cos \phi}{\Delta}.$$

## CHAPTER II.

### ELLIPTIC FUNCTIONS.

LET 
$$u = \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}.$$

$\phi^*$  is called the *amplitude* corresponding to the *argument*  $u$ , and is written

$$\phi = \text{am}(u, k) = \text{am } u.$$

The quantity  $k$  is called the *modulus*, and the expression  $\sqrt{1 - k^2 \sin^2 \phi}$  is written\*

$$\sqrt{1 - k^2 \sin^2 \phi} = \Delta \text{am } u = \Delta \phi,$$

and is called the *delta function* of the amplitude of  $u$ , or *delta of  $\phi$* , or simply *delta  $\phi$* .

$u$  can be written

$$u = F(k, \phi).$$

The following abbreviations are used:

$$\sin \phi = \sin \text{am } u = \text{sn}^\dagger u;$$

$$\cos \phi = \cos \text{am } u = \text{cn}^\dagger u;$$

$$\Delta \phi = \Delta \text{am } u = \text{dn}^\dagger u = \Delta u;$$

$$\tan \phi = \tan \text{am } u = \text{tn } u.$$

---

\*Legendre.

†Gudermann, in his "Theorie der Modularfunctionen": Crelle's Journal, Bd. 18.

Let  $\phi$  and  $\psi$  be any two arbitrary angles, and put

$$\phi = \text{am } u;$$


$$\psi = \text{am } v.$$

In the spherical triangle  $ABC$  we have from Trigonometry,  $\mu$  and  $C$  being constant,

$$\frac{d\phi}{\cos B} + \frac{d\psi}{\cos A} = 0.$$

Since  $C$  and  $\mu$  are constant, denoting by  $k$  an arbitrary constant, we have

$$(1) \quad \frac{\sin C}{\sin \mu} = k.$$



But

$$\sin A = \sin \psi \frac{\sin B}{\sin \phi} = \sin \psi \frac{\sin C}{\sin \mu} = k \sin \psi.$$

Whence

$$\cos A = \sqrt{1 - \sin^2 A} = \sqrt{1 - k^2 \sin^2 \psi}.$$

In the same manner

$$\cos B = \sqrt{1 - \sin^2 B} = \sqrt{1 - k^2 \sin^2 \phi}.$$

Substituting these values, we get

$$(2) \quad \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} + \frac{d\psi}{\sqrt{1 - k^2 \sin^2 \psi}} = 0.$$



Integrating this, there results

$$(3) \quad \int_0^\phi \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} + \int_0^\psi \frac{d\psi}{\sqrt{1-k^2 \sin^2 \psi}} = \text{const.}$$

When  $\phi = 0$ , we have  $\psi = \mu$ , and therefore the constant must be of the form

$$\int_0^\mu \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}},$$

whence

$$(4) \quad \int_0^\phi \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} + \int_0^\psi \frac{d\psi}{\sqrt{1-k^2 \sin^2 \psi}} = \int_0^\mu \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}},$$

or

$$u + v = m;$$


and evidently the amplitudes  $\phi$ ,  $\psi$ , and  $\mu$  can be considered as the three sides of a spherical triangle, and the relations between the sides of this spherical triangle will be the same as those between  $\phi$ ,  $\psi$ , and  $\mu$ .

But the sides of this triangle have imposed upon them the condition

$$\frac{\sin C}{\sin \mu} = k;$$


and since  $k < 1$ , we must have  $\mu > C$ , which requires that one of the angles of the triangle shall be obtuse and the other two acute.

In the figure, let  $C$  be an acute angle of the triangle  $ABC$ , and  $PQ$  the equatorial great circle of which  $C$  is the pole.



The arc  $PQ$  will be the measure of the angle  $C$ .

Let  $AG$  and  $AH$  be the arcs of two great circles perpendicular respectively to  $CQ$  and  $CP$ . They will of course be shorter than  $PQ$ . Hence  $AB = \mu$  must intersect  $CQ$  in points between  $CG$  and  $HQ$ , since  $\mu > (C = PQ)$ . In any case either  $A$  or  $B$  will be obtuse according as  $B$  falls between  $QH$  or  $CG$  respectively; and the other angle will be acute.



In the case where  $C$  is an obtuse angle, it will be easily seen that the angle at  $A$  must be acute, since the great circle  $AD$ , perpendicular to  $CP$ , intersects  $PQ$  in  $D$ ,  $PD$  being a quadrant. The same remarks apply to the angle  $B$ . Hence, in either case, one of the angles of the triangle is obtuse and the other two are acute, as a result of the condition

$$\frac{\sin C}{\sin \mu} = k < 1.$$

From Trigonometry we have

$$\cos \mu = \cos \phi \cos \psi + \sin \phi \sin \psi \cos C;$$

and since the angle  $C$  is obtuse,

$$\cos C = -\sqrt{1 - \sin^2 C} = -\sqrt{1 - k^2 \sin^2 \mu},$$

and

$$(5) \quad \cos \mu = \cos \phi \cos \psi - \sin \phi \sin \psi \sqrt{1 - k^2 \sin^2 \mu},$$

the relation sought.

The spherical triangle likewise gives the following relations between the sides:

$$(5)^* \quad \begin{cases} \cos \phi = \cos \mu \cos \psi + \sin \mu \sin \psi \sqrt{1 - k^2 \sin^2 \phi}; \\ \cos \psi = \cos \mu \cos \phi + \sin \mu \sin \phi \sqrt{1 - k^2 \sin^2 \psi}. \end{cases}$$

These give, by eliminating  $\cos \mu$ ,

$$\sin \mu = \frac{\cos^2 \psi - \cos^2 \phi}{\sin \phi \cos \psi \Delta \psi - \sin \psi \cos \phi \Delta \phi};$$

which, after multiplying by the sum of the terms in the denominator and substituting  $\cos^2 = 1 - \sin^2$ , can be written

$$\sin \mu = \frac{(\sin^2 \phi - \sin^2 \psi)(\sin \phi \cos \psi \Delta \psi + \sin \psi \cos \phi \Delta \phi)}{\sin^2 \phi \cos^2 \psi \Delta^2 \psi - \sin^2 \psi \cos^2 \phi \Delta^2 \phi}.$$

Since the denominator can be written

$$(6) \quad \begin{aligned} & (\sin^2 \phi - \sin^2 \psi)(1 - k^2 \sin^2 \phi \sin^2 \psi), \\ \sin \mu &= \frac{\sin \phi \cos \psi \Delta \psi + \sin \psi \cos \phi \Delta \phi}{1 - k^2 \sin^2 \phi \sin^2 \psi}. \end{aligned}$$

In a similar manner we get

$$(6)^* \quad \begin{cases} \cos \mu = \frac{\cos \phi \cos \psi - \sin \phi \sin \psi \Delta \phi \Delta \psi}{1 - k^2 \sin^2 \phi \sin^2 \psi}; \\ \Delta \mu = \frac{\Delta \phi \Delta \psi - k^2 \sin \phi \sin \psi \cos \phi \cos \psi}{1 - k^2 \sin^2 \phi \sin^2 \psi}. \end{cases}$$

These equations can also be written as follows:

$$(7) \quad \begin{cases} \sin \operatorname{am}(u \pm v) = \frac{\sin \operatorname{am} u \cos \operatorname{am} v \Delta \operatorname{am} v \pm \sin \operatorname{am} v \cos \operatorname{am} u \Delta \operatorname{am} u}{1 - k^2 \sin^2 \operatorname{am} u \sin^2 \operatorname{am} v}; \\ \cos \operatorname{am}(u \pm v) = \frac{\cos \operatorname{am} u \cos \operatorname{am} v \mp \sin \operatorname{am} u \sin \operatorname{am} v \Delta \operatorname{am} u \Delta \operatorname{am} v}{1 - k^2 \sin^2 \operatorname{am} u \sin^2 \operatorname{am} v}; \\ \Delta \operatorname{am}(u \pm v) = \frac{\Delta \operatorname{am} u \Delta \operatorname{am} v \mp k^2 \sin \operatorname{am} u \sin \operatorname{am} v \cos \operatorname{am} u \cos \operatorname{am} v}{1 - k^2 \sin^2 \operatorname{am} u \sin^2 \operatorname{am} v}; \end{cases}$$

or

$$(8) \quad \begin{cases} \operatorname{sn}(u \pm v) = \frac{\operatorname{sn} u \operatorname{cn} v \operatorname{dn} v \pm \operatorname{sn} v \operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}; \\ \operatorname{cn}(u \pm v) = \frac{\operatorname{cn} u \operatorname{cn} v \mp \operatorname{sn} u \operatorname{sn} v \operatorname{dn} u \operatorname{dn} v}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}; \\ \operatorname{dn}(u \pm v) = \frac{\operatorname{dn} u \operatorname{dn} v \mp k^2 \operatorname{sn} u \operatorname{sn} v \operatorname{cn} u \operatorname{cn} v}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}. \end{cases}$$

Making  $u = v$ , we get from the upper sign

$$(9) \quad \begin{cases} \operatorname{sn} 2u = \frac{2 \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^4 u}; \\ \operatorname{cn} 2u = \frac{\operatorname{cn}^2 u - \operatorname{sn}^2 u \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u} = \frac{1 - 2 \operatorname{sn}^2 u + k^2 \operatorname{sn}^4 u}{1 - k^2 \operatorname{sn}^4 u}; \\ \operatorname{dn} 2u = \frac{\operatorname{dn}^2 u - k^2 \operatorname{sn}^2 u \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u} = \frac{1 - 2k^2 \operatorname{sn}^2 u + k^2 \operatorname{sn}^4 u}{1 - k^2 \operatorname{sn}^4 u}. \end{cases}$$

From these

$$(10) \quad \begin{cases} 1 - \operatorname{cn} 2u = \frac{2 \operatorname{cn}^2 u \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \\ 1 + \operatorname{cn} 2u = \frac{2 \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \\ 1 - \operatorname{dn} u = \frac{2k^2 \operatorname{sn}^2 u \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \\ 1 + \operatorname{dn} u = \frac{2 \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \end{cases}$$

and therefore

$$(11) \quad \begin{cases} \operatorname{sn}^2 u = \frac{1 - \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}; \\ \operatorname{cn}^2 u = \frac{\operatorname{dn} 2u + \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}; \\ \operatorname{dn}^2 u = \frac{1 - k^2 + \operatorname{dn} 2u + k^2 \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}; \end{cases}$$

and by analogy

$$(12) \quad \begin{cases} \operatorname{sn} \frac{u}{2} = \sqrt{\frac{1 - \operatorname{cn} u}{1 + \operatorname{dn} u}}; \\ \operatorname{cn} \frac{u}{2} = \sqrt{\frac{\operatorname{cn} u + \operatorname{dn} u}{1 + \operatorname{dn} u}}; \\ \operatorname{dn} \frac{u}{2} = \sqrt{\frac{1 - k^2 + \operatorname{dn} u + k^2 \operatorname{cn} u}{1 + \operatorname{dn} u}}. \end{cases}$$

In [equations \(7\)](#) making  $u = v$ , and taking the lower sign, we have

$$(13) \quad \begin{cases} \operatorname{sn} 0 = 0; \\ \operatorname{cn} 0 = 1; \\ \operatorname{dn} 0 = 1. \end{cases}$$

Likewise, we get by making  $u = 0$ ,

$$(14) \quad \begin{cases} \operatorname{sn}(-u) = -\operatorname{sn} u; \\ \operatorname{cn}(-u) = +\operatorname{cn} u; \\ \operatorname{dn}(-u) = \operatorname{dn} u. \end{cases}$$

# CHAPTER III.

## PERIODICITY OF THE FUNCTIONS.

WHEN the elliptic integral

$$\int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}$$

has for its amplitude  $\frac{\pi}{2}$ , it is called, following the notation of Legendre, the *complete* function, and is indicated by  $K$ , thus:

$$K = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}.$$

When  $k$  becomes the complementary modulus,  $k'$ , (see [eq. \(4\)](#), Chap. IV,) the corresponding complete function is indicated by  $K'$ , thus:

$$K' = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1 - k'^2 \sin^2 \phi}}.$$

From these, evidently,

$$\text{am}(K, k) = \frac{\pi}{2}, \quad \text{am}(K', k') = \frac{\pi}{2}.$$

$$(1) \quad \begin{cases} \text{sn}(K, k) = 1; \\ \text{cn}(K, k) = 0; \\ \text{dn}(K, k) = k'. \end{cases}$$

From eqs. (7), (8), and (9), Chap. II, we have, by the substitution of the values of  $\text{sn}(K) = 1$ ,  $\text{cn}(K) = 0$ ,  $\text{dn}(K) = k'$ ,

$$(2) \quad \begin{cases} \text{sn } 2K = 0; \\ \text{cn } 2K = -1; \\ \text{dn } 2K = 1. \end{cases}$$

These equations, by means of (1), (2), and (3) of Chap. II, give

$$(3) \quad \begin{cases} \text{sn}(u + 2K) = -\text{sn } u; \\ \text{cn}(u + 2K) = -\text{cn } u; \\ \text{dn}(u + 2K) = \text{dn } u; \end{cases}$$

and these, by changing  $u$  into  $u + 2K$ , give

$$(4) \quad \begin{cases} \text{sn}(u + 4K) = \text{sn } u; \\ \text{cn}(u + 4K) = \text{cn } u; \\ \text{dn}(u + 4K) = \text{dn } u. \end{cases}$$

From these equations it is seen that the elliptic functions  $\text{sn}$ ,  $\text{cn}$ ,  $\text{dn}$ , are periodic functions having for their period  $4K$ . Unlike the period of trigonometric functions, this period is not a fixed one, but depends upon the value of  $k$ , the modulus.

From the Integral Calculus we have

$$\begin{aligned} \int_0^{n\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} &= \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} + \int_{\frac{\pi}{2}}^{\pi} \frac{d\phi}{\Delta\phi} + \int_{\pi}^{\frac{3\pi}{2}} \frac{d\phi}{\Delta\phi} + \cdots + \int_{(n-1)\frac{\pi}{2}}^{n\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} \\ &= n \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} = nK; \end{aligned}$$

from which we see that

$$n\frac{\pi}{2} = \text{am}(nK);$$



or, since  $\frac{\pi}{2} = \text{am } K$ ,

$$\text{am}(nK) = n \cdot \text{am } K,$$

and

$$n\pi = \text{am}(2nK),$$

and also

$$n\pi = 2n \text{ am } K.$$

In the case of an Elliptic Integral with the arbitrary angle  $\alpha$ , we can put

$$\alpha = n\pi \pm \beta,$$

where  $\beta$  is an angle between 0 and  $\frac{\pi}{2}$ , the upper or the lower sign being taken according as  $\frac{\pi}{2}$  is contained in  $\alpha$  an even or an uneven number of times.

In the first case we have

$$\int_0^{n\pi+\beta} \frac{d\phi}{\Delta\phi} = \int_0^{n\pi} \frac{d\phi}{\Delta\phi} + \int_{n\pi}^{n\pi+\beta} \frac{d\phi}{\Delta\phi};$$

or, putting  $\phi_1 = \phi - n\pi$ ,

$$\int_0^{n\pi+\beta} \frac{d\phi}{\Delta\phi} = 2nK + \int_0^\beta \frac{d\phi_1}{\Delta\phi_1}.$$

In the second case

$$\int_0^{n\pi-\beta} \frac{d\phi}{\Delta\phi} = \int_0^{n\pi} \frac{d\phi}{\Delta\phi} - \int_{n\pi-\beta}^{n\pi} \frac{d\phi}{\Delta\phi};$$

or, putting  $\phi_1 = n\pi - \phi$ ,

$$\int_0^{n\pi-\beta} \frac{d\phi}{\Delta\phi} = 2nK - \int_0^\beta \frac{d\phi_1}{\Delta\phi_1};$$

or in either case,

$$\int_0^{n\pi \pm \beta} \frac{d\phi}{\Delta\phi} = 2nK \pm \int_0^\beta \frac{d\phi_1}{\Delta\phi_1}.$$

Thus we see that the Integral with the general amplitude  $\alpha$  can be made to depend upon the complete integral  $K$  and an Integral whose amplitude lies between 0 and  $\frac{\pi}{2}$ .

Put now

$$\int_0^\beta \frac{d\phi_1}{\Delta\phi_1} = u, \quad \beta = \text{am } u.$$

This gives

$$\int_0^{n\pi \pm \beta} \frac{d\phi}{\Delta\phi} = 2nK \pm u,$$

or  $\text{am}(2nK \pm u) = n\pi \pm \beta$

(5)  $\qquad \qquad \qquad = n\pi \pm \text{am } u$

(6)  $\qquad \qquad \qquad = 2n \cdot \text{am } K \pm \text{am } u;$

or, since

$$\text{am}(-z) = -\text{am } z,$$

$$\text{am}(u \pm 2nK) = \text{am } u \pm n\pi$$

$$= \text{am } u \pm 2n \cdot \text{am } K.$$

Taking the sine and cosine of both sides, we have

$$\text{sn}(u + 2nK) = \pm \text{sn } u;$$

$$\text{cn}(u + 2nK) = \pm \text{cn } u;$$

the upper or the lower sign being taken according as  $n$  is even or odd. By giving the proper values to  $n$  we can get the same results as in equations (3) and (4).

Putting  $n = 1$  in eq. (5), we have

$$(7) \quad \begin{aligned} \operatorname{sn}(2K - u) &= \sin \pi \operatorname{cn} u - \cos \pi \operatorname{sn} u \\ &= \operatorname{sn} u. \end{aligned}$$

Elliptic functions also have an imaginary period. In order to show this we will, in the integral

$$\int_0^\phi \frac{d\phi}{\Delta\phi'}$$

assume the amplitude as imaginary. Put

$$\sin \phi = i \tan \psi.$$

From this we get

$$(8) \quad \left\{ \begin{aligned} \cos \phi &= \frac{1}{\cos \psi}; \\ \Delta\phi &= \frac{\sqrt{1 - k'^2 \sin^2 \psi}}{\cos \psi} = \frac{\Delta(\psi, k')}{\cos \psi}; \\ d\phi &= i \frac{d\psi}{\cos \psi}. \end{aligned} \right.$$

From these, since  $\phi$  and  $\psi$  vanish simultaneously, we easily get

$$\int_0^\phi \frac{d\phi}{\Delta\phi} = i \int_0^\psi \frac{d\psi}{\Delta(\psi, k')}.$$

Put

$$\int_0^\psi \frac{d\psi}{\Delta(\psi, k')} = u \quad \text{and} \quad \psi = \operatorname{am}(u, k'),$$

whence

$$\int_0^\phi \frac{d\phi}{\Delta\phi} = iu \quad \text{and} \quad \phi = \text{am}(iu);$$

and these substituted in Eq. (8) give

$$(9) \quad \begin{cases} \text{sn } iu = i \text{tn}(u, k'); \\ \text{cn } iu = \frac{1}{\text{cn}(u, k')}; \\ \text{dn } iu = \frac{\text{dn}(u, k')}{\text{cn}(u, k')}. \end{cases}$$

By assuming

$$\int_0^\psi \frac{d\psi}{\Delta(\psi, k')} = iu \quad \text{and} \quad \int_0^\phi \frac{d\phi}{\Delta\phi} = -u,$$

we get

$$\begin{aligned} \text{sn}(-u) &= i \text{tn}(iu, k'), \\ \text{cn}(-u) &= \frac{1}{\text{cn}(iu, k')}, \\ \text{dn}(-u) &= \frac{\text{dn}(iu, k')}{\text{cn}(iu, k')}; \end{aligned}$$

or, from eq. (14), Chap. II,

$$(10) \quad \begin{cases} \text{sn } u = -i \text{tn}(iu, k'); \\ \text{cn } u = \frac{1}{\text{cn}(iu, k')}; \\ \text{dn } u = \frac{\text{dn}(iu, k')}{\text{cn}(iu, k')}. \end{cases}$$

From eqs. (7), Chap. II, making  $v = K$ , we get, since  $\operatorname{sn} K = 1$ ,  $\operatorname{cn} K = 0$ ,  $\operatorname{dn} K = k'$ ,

$$(11) \quad \begin{cases} \operatorname{sn}(u \pm K) = \pm \frac{\operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^2 u} = \pm \frac{\operatorname{cn} u}{\operatorname{dn} u}; \\ \operatorname{cn}(u \pm K) = \mp \frac{\operatorname{sn} u \operatorname{dn} u k'}{\operatorname{dn}^2 u} = \mp \frac{k' \operatorname{sn} u}{\operatorname{dn} u}; \\ \operatorname{dn}(u \pm K) = + \frac{k'}{\operatorname{dn} u}. \end{cases}$$

In these equations, changing  $u$  into  $iu$ , we get, by means of eqs. (9),

$$(12) \quad \begin{cases} \operatorname{sn}(iu \pm K) = \pm \frac{1}{\operatorname{dn}(u, k')}; \\ \operatorname{cn}(iu \pm K) = \mp \frac{ik' \operatorname{sn}(u, k')}{\operatorname{dn}(u, k')}; \\ \operatorname{dn}(iu \pm K) = + \frac{k' \operatorname{cn}(u, k')}{\operatorname{dn}(u, k')}. \end{cases}$$

Putting now in eqs. (9)  $u \pm K'$  instead of  $u$ , and making use of eqs. (10), and interchanging  $k$  and  $k'$ , we have

$$(13) \quad \begin{cases} \operatorname{sn}(iu \pm iK') = - \frac{i \operatorname{cn}(u, k')}{k \operatorname{sn}(u, k')}; \\ \operatorname{cn}(iu \pm iK') = \mp \frac{\operatorname{dn}(u, k')}{k \operatorname{sn}(u, k')}; \\ \operatorname{dn}(iu \pm iK') = \mp \frac{1}{\operatorname{sn}(u, k')}. \end{cases}$$

Substituting in these  $-iu$  in place of  $u$ , we get, by means of eqs. (9)

and eqs. (14) of Chap. II,

$$(14) \quad \begin{cases} \operatorname{sn}(u \pm iK') = \frac{1}{k \operatorname{sn} u}; \\ \operatorname{cn}(u \pm iK') = \mp \frac{i \operatorname{dn} u}{k \operatorname{sn} u}; \\ \operatorname{dn}(u \pm iK') = \mp i \cot \operatorname{am} u. \end{cases}$$

In these equations, putting  $u + K$  in place of  $u$ , we get

$$(15) \quad \begin{cases} \operatorname{sn}(u + K \pm iK') = + \frac{\operatorname{dn} u}{k \operatorname{cn} u}; \\ \operatorname{cn}(u + K \pm iK') = \mp \frac{ik'}{k \operatorname{cn} u}; \\ \operatorname{dn}(u + K \pm iK') = \pm ik' \operatorname{tn} u. \end{cases}$$

Whence for  $u = 0$  we get

$$(16) \quad \begin{cases} \operatorname{sn}(K \pm iK') = \frac{1}{k}; \\ \operatorname{cn}(K \pm iK') = \mp \frac{ik'}{k}; \\ \operatorname{dn}(K \pm iK') = 0. \end{cases}$$

If in eqs. (14) we put  $u = 0$ , we see that as  $u$  approaches zero, the expressions

$$\operatorname{sn}(\pm iK'), \quad \operatorname{cn}(\pm iK'), \quad \operatorname{dn}(\pm iK')$$

approach infinity.

We see from what has preceded that Elliptic Functions have two periods, one a real period, and one an imaginary period.

In the former characteristic they resemble Trigonometric Functions, and in the latter Logarithmic Functions.

On account of these two periods they are often called Doubly Periodic Functions. Some authors make this double periodicity the starting-point of their investigations. This method of investigation gives some very beautiful results and processes, but not of a kind adapted for an elementary work.


It will be noticed that the Elliptic Functions  $\operatorname{sn} u$ ,  $\operatorname{cn} u$ , and  $\operatorname{dn} u$  have a very close analogy to trigonometric functions, in which, however, the independent variable  $u$  is not an angle, as it is in the case of trigonometric functions.

Like Trigonometric Functions, these Elliptic Functions can be arranged in tables. These tables, however, require a double argument, viz.,  $u$  and  $k$ . In [Chap. IX](#) these functions are developed into series, from which their values may be computed and tables formed.

No complete tables have yet been published, though they are in process of computation.

# CHAPTER IV.

## LANDEN'S TRANSFORMATION



LET  $AB$  be the diameter of a circle, with the centre at  $O$ , the radius  $AO = r$ , and  $C$  a fixed point situated upon  $OB$ , and  $OC = k_0 r$ . Denote the angle  $PBC$  by  $\phi$ , and the angle  $PCO$  by  $\phi_1$ . Let  $P'$  be a point indefinitely near to  $P$ .

Then

$$\frac{PP'}{PC} = \frac{\sin PCP'}{\sin PP'C} = \frac{\sin PCP'}{\cos OP'C}.$$

But  $PP' = 2r d\phi$ , and  $\sin PCP' = PCP' = d\phi_1$ ; therefore

$$\frac{2r d\phi}{PC} = \frac{d\phi_1}{\cos OP'C}.$$

But

$$\begin{aligned} \overline{PC}^2 &= r^2 + r^2 k_0^2 + 2r^2 k_0 \cos 2\phi \\ &= (r + rk_0)^2 \cos^2 \phi + (r - rk_0)^2 \sin^2 \phi; \\ \text{also } r^2 \cos^2 OP'C &= r^2 - r^2 \sin^2 OP'C \\ &= r^2 - r^2 k_0^2 \sin^2 \phi_1. \end{aligned}$$

Therefore

$$\frac{2 d\phi}{\sqrt{(r + rk_0)^2 \cos^2 \phi + (r - rk_0)^2 \sin^2 \phi}} = \frac{d\phi_1}{\sqrt{r^2 - r^2 k_0^2 \sin^2 \phi_1}},$$

which can be written

$$\frac{2}{r + rk_0} \frac{d\phi}{\sqrt{1 - \frac{4k_0^2 r^2}{(r + rk_0)^2} \sin^2 \phi}} = \frac{1}{r} \frac{d\phi_1}{\sqrt{1 - k_0^2 \sin^2 \phi_1}},$$



Putting

$$(1) \quad \frac{4k_0r^2}{(r + rk_0)^2} = \frac{4k_0}{(1 + k_0)^2} = k^2,$$

we have

$$(2) \quad \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = \frac{1 + k_0}{2} \int_0^{\phi_1} \frac{d\phi_1}{\sqrt{1 - k_0^2 \sin^2 \phi_1}};$$

no constant being added because  $\phi$  and  $\phi_1$  vanish simultaneously;  $\phi$  and  $\phi_1$  being connected by the equation

$$(3) \quad \frac{\sin OPC}{\sin OCP} = \frac{\sin(2\phi - \phi_1)}{\sin \phi_1} = \frac{rk_0}{r} = k_0.$$

From the value of  $k^2$  we have

$$(4) \quad 1 - k^2 = k'^2 = \frac{(1 - k_0)^2}{(1 + k_0)^2},$$

and therefore

$$(5) \quad k_0 = \frac{1 - k'}{1 + k'}.$$

$k'$  is called the *complementary modulus*, and is evidently the minimum value of  $\Delta\phi$ , the value of  $\Delta\phi$  when  $\phi = 90^\circ$ :

$$\sqrt{1 - k^2} = k'.$$

From [eq. \(1\)](#) we evidently have  $k > k_0$ , for, putting [eq. \(1\)](#) in the form

$$\frac{k^2}{k_0^2} = \frac{4}{k_0 + 2k_0^2 + k_0^3},$$

we see that if  $k_0 = 1$ , then  $k = k_0$ , but as  $k_0 < 1$ , always, as is evident from the figure,  $k$  must be greater than  $k_0$ .

It is also evident, from the figure, that  $\phi_1 > \phi$ . Or it may be deduced directly from [eq. \(3\)](#).

Since  $k < 1$ , we can write

$$k = \sin \theta, \quad k' = \sqrt{1 - k^2} = \cos \theta.$$

Substituting in [eq. \(5\)](#), we have

$$k_0 = \frac{1 - k'}{1 + k'} = \tan^2 \frac{1}{2}\theta,$$

and we can write

$$k_0 = \sin \theta_0, \quad k'_1 = \sqrt{1 - k_0^2} = \cos \theta_0.$$

From [eq. \(5\)](#) we have

$$1 + k_0 = \frac{2}{1 + k'}.$$

Substituting the value of  $k_0$  in that for  $k'_1$ , we get

$$k'_1 = \frac{2\sqrt{k'}}{1 + k'}.$$

We also have

$$\begin{aligned} 2\phi - \phi_1 &= \phi - (\phi_1 - \phi) \\ \phi_1 &= \phi + (\phi_1 - \phi), \end{aligned}$$

and, [eq. \(3\)](#),

$$\operatorname{sn}(2\phi - \phi_1) = k_0 \sin \phi_1,$$

becomes

$$\begin{aligned} \sin \phi \cos(\phi_1 - \phi) - \cos \phi \sin(\phi_1 - \phi) \\ = k_0 \sin \phi \cos(\phi_1 - \phi) + k_0 \cos \phi \sin(\phi_1 - \phi), \end{aligned}$$

or

$$\tan \phi - \tan(\phi_1 - \phi) = k_0 \tan \phi + k_0 \tan(\phi_1 - \phi),$$

or

$$\begin{aligned} \tan(\phi_1 - \phi) &= \frac{1 - k_0}{1 + k_0} \tan \phi \\ &= k' \tan \phi. \end{aligned}$$

Collecting these results, we have

$$(6) \quad k = \frac{2\sqrt{k_0}}{1 + k_0} = \sin \theta;$$

$$(7) \quad k_0 = \frac{1 - k'}{1 + k'} = \sin \theta_0 = \tan^2 \frac{1}{2}\theta;$$

$$(8) \quad k'_1 = \frac{2\sqrt{k'}}{1 + k'} = \cos \theta_0;$$

$$(9) \quad k' = \frac{1 - k_0}{1 + k_0} = \cos \theta;$$

$$(10) \quad 1 + k_0 = \frac{2}{1 + k'} = \frac{2\sqrt{k_0}}{k} = \frac{k'_1}{\sqrt{k'}} = \frac{1}{\cos^2 \frac{1}{2}\theta};$$

$$(11) \quad \sin(2\phi - \phi_1) = k_0 \sin \phi_1;$$

$$(12) \quad \tan(\phi_1 - \phi) = k' \tan \phi;$$

$$(13) \quad \int_0^\phi \frac{d\phi}{\Delta(k, \phi)} = \frac{1+k_0}{2} \int_0^{\phi_1} \frac{d\phi_1}{\Delta(k_0, \phi_1)};$$

$$(14) \quad k = \sqrt{1-k'^2}, \quad k' = \sqrt{1-k^2}.$$

Upon examination it will easily appear that  $k$  and  $k_0$ , and  $\theta$  and  $\theta_0$ , are the first two terms of a decreasing series of moduli and angles;  $k'$  and  $k'_1$ , and  $\phi$  and  $\phi_1$ , of an increasing series; the law connecting the different terms of the series being deduced from eqs. (6) to (12).

By repeated applications of these equations we would get the following series of moduli and amplitudes:

$k_{0n} = 0_{(n=\infty)}$	$k'_n = 1_{(n=\infty)}$	$\phi_n$
$\vdots$	$\vdots$	$\vdots$
$k_{00}$	$k'_2$	$\phi_2$
$k_0$	$k'_1$	$\phi_1$
$k$	$k'$	$\phi$

The upper limit of the one series of moduli is 1, and the lower limit of the other series is 0, as is indicated.  $k$  and  $k'$ , which are bound by the relation  $k^2 + k'^2 = 1$ , are called the *primitives* of the series.

NOTE.— It will be noticed that the successive terms of a decreasing series are indicated by the sub-accents 0, 00, 03, 04, . . . 0n; and the successive terms of an increasing series by the sub-accents 1, 2, 3, . . . n.

Again, by application of these equations, we can form a new series running up from  $k$ , viz.,  $k_1, k_2, k_3, \dots k_n = 1_{(n=\infty)}$ ; and also a new series running down from  $k'$ , viz.,  $k'_0, k'_{00}, \dots k'_{0n} = 0_{(n=\infty)}$ . So also with  $\phi$ .





From eqs. (15) and (10) we get

$$F(k, \phi) = \sqrt{\frac{k'_1 k'_2 k'_3 \cdots k'_n}{k'}} \cdot \frac{F(k_{0n}, \phi_n)}{2^n}.$$

And this with equations (8) and (9) gives

$$(17) \quad F(k, \phi) = \sqrt{\frac{\cos \theta_0 \cos \theta_{00} \cdots \cos^2 \theta_{0n}}{\cos \theta}} \cdot \frac{F(k_{0n}, \phi_n)}{2^n}.$$

Applying equation (13) to  $(k_1, \phi_0)$ ,  $(k_2, \phi_{00})$ , etc., we get

$$F(k_1, \phi_0) = \frac{1+k}{2} F(k, \phi), \text{ etc.};$$

but since, eq. (10),

$$\frac{1+k}{2} = \frac{1}{1+k'_0}, \text{ etc.},$$

these become

$$F(k, \phi) = (1+k'_0)F(k_1, \phi_0);$$

$$F(k_1, \phi_0) = (1+k'_{00})F(k_2, \phi_{00});$$

· · · · ·

$$F(k_{n-1}, \phi_{0(n-1)}) = (1+k'_{0n})F(k_n, \phi_{0n});$$

whence

$$(18) \quad F(k, \phi) = (1+k'_0)(1+k'_{00}) \cdots (1+k'_{0n})F(k_n, \phi_{0n}),$$

in which  $k'_0$ ,  $k'_{00}$ , etc.,  $k_1$ ,  $k_2$ , etc.,  $\phi_0$ ,  $\phi_{00}$ , etc., are determined as follows:

Let  $k = \sin \theta,$   
 $k_1 = \sin \theta_1.$

From eq. (10),

$$k_1 = \frac{2\sqrt{k}}{1+k} \quad \text{or} \quad \sin \theta_1 = \frac{2\sqrt{\sin \theta}}{1 + \sin \theta}.$$

Solving this equation for  $\sin \theta$ , we get

$$\sin \theta = \tan^2 \frac{1}{2} \theta_1.$$

Hence we can write

$$(18_1) \quad \left\{ \begin{array}{l} k = \sin \theta = \tan^2 \frac{1}{2} \theta_1; \\ k_1 = \sin \theta_1 = \tan^2 \frac{1}{2} \theta_2; \\ \dots \dots \dots \\ k_n = \sin \theta_n. \end{array} \right.$$

From equation (12) we get

$$(18_2) \quad \left\{ \begin{array}{l} \sin(2\phi_0 - \phi) = k \sin \phi;^* \\ \sin(2\phi_{00} - \phi_0) = k_1 \sin \phi_0; \\ \dots \dots \dots \\ \sin(2\phi_{0n} - \phi_{0(n-1)}) = k_{n-1} \sin \phi_{0(n-1)}. \end{array} \right.$$

---

\*When  $\sin \phi = 1$  nearly,  $\phi$  is best determined as follows: From eq. (12) we have

$$\begin{aligned} \tan(\phi - \phi_0) &= k'_0 \tan \phi_0 \\ &= k'_0 \tan \phi \text{ nearly;} \end{aligned}$$



---

whence

$$\phi - \phi_0 = Rk'_0 \tan \phi \text{ nearly,}$$

$R$  being the radian in seconds, viz.  $206264'' .806$ , and  $\log R = 5.3144251$ .

Substituting the approximate value of  $\phi_0$ , we can get a new approximation.

*Example.*       $\phi_0 = 82^\circ 30'$        $k'_{00} = \log^{-1} 5.8757219$

$\tan 82^\circ 30'$	10.8805709	
$k'_{00}$	5.8757219	
$R$	5.3144251	
	2.0707179	$117'' .684 = 1'.9614$

$$\begin{aligned} \phi_0 - \phi_{00} &= 1'.9614 \\ \phi_{00} &= 82^\circ 28'.0386 && \text{1st approximation.} \end{aligned}$$

This value gives

$$\begin{aligned} \phi_0 - \phi_{00} &= 117'' .1675 = 1'.95279 \\ \therefore \phi_{00} &= 82^\circ 28'.04721 && \text{2d approximation.} \end{aligned}$$

This value gives

$$\begin{aligned} \phi_0 - \phi_{00} &= 117'' .1698 = 1'.95283 \\ \phi_{00} &= 82^\circ 28'.04717 && \text{3d approximation.} \end{aligned}$$

To determine  $k'_0, k'_{00}$ , etc., we have

$$(18_3) \quad \left\{ \begin{array}{ll} k' = \sin \eta, & k = \cos \eta; \\ k'_0 = \frac{1-k}{1+k} = \tan^2 \frac{1}{2} \eta = \sin \eta_0, & k_1 = \cos \eta_0; \\ k'_{00} = \tan^2 \frac{1}{2} \eta_0 = \sin \eta_{00}, & k_2 = \cos \eta_{00}; \\ \text{etc.} & \text{etc.} \quad \text{etc.} \end{array} \right.$$

Or, since  $1 + k'_0 = \frac{1}{\cos^2 \frac{1}{2} \eta}$ ,  $1 + k'_{00} = \frac{1}{\cos^2 \frac{1}{2} \eta_0}$ , etc., we can put eq. (18) in the following form:

$$(19) \quad F(k, \phi) = \frac{1}{\cos^2 \frac{1}{2} \eta \cos^2 \frac{1}{2} \eta_0 \cdots \cos^2 \frac{1}{2} \eta_{0n}} F(k_n, \phi_{0n}).$$

From equation (13) we have

$$(19)^* \quad F(k_1, \phi_0) = \frac{1+k}{2} F(k, \phi),$$

whence

$$F(k, \phi) = \frac{2}{1+k} F(k_1, \phi_0).$$

By repeated applications this gives, after combining,

$$(20) \quad \begin{aligned} F(k, \phi) &= \frac{2}{1+k} \cdot \frac{2}{1+k_1} \cdots \frac{2}{1+k_{n-1}} \cdot F(k_n, \phi_{0n}) \\ &= \frac{k_1}{\sqrt{k}} \cdot \frac{k_2}{\sqrt{k_1}} \cdots \frac{k_n}{\sqrt{k_{n-1}}} \cdot F(k_n, \phi_{0n}); \\ &= \sqrt{\frac{k_1 k_2 \cdots k_n^2}{k}} \cdot F(k_n, \phi_{0n}); \end{aligned}$$

$k_1, k_2$ , etc., being determined by repeated applications of

$$k_1 = \frac{2\sqrt{k}}{1+k'}$$

or by [equations \(18<sub>1</sub>\)](#).

In [equation \(19\)\\*](#) let us change  $k_1$  and  $\phi_0$  into  $k'$  and  $\phi$  respectively, so that the first member may have for its complete function

$$K' = F(k', \phi).$$

Upon examination of [eq. \(19\)\\*](#) we see that the modulus in the second member must be the one next less than the one in the first member, that is,  $k'_0$ ; and likewise that the amplitude must be the one next greater than the amplitude in the first member, viz.,  $\phi_1$ ; hence we get

$$F(k', \phi) = \frac{1+k'_0}{2} F(k'_0, \phi_1).$$

Indicating the complete functions by  $K'$  and  $K'_0$ , we have, since  $\phi = \frac{\pi}{2}$  when  $\phi_1 = \pi$  (see [Chap. V](#)),

$$K' = (1+k'_0)K'_0;$$

and in the same manner,

$$K'_0 = (1+k'_{00})K'_{00},$$

$$K'_{00} = (1+k'_{03})K'_{03},$$

· · · · ·

$$K'_{0(n-1)} = (1+k'_{0n})K'_{0n};$$

whence

$$K' = (1+k'_0)(1+k'_{00}) \cdots (1+k'_{0n})K'_{0n}.$$

Since

$$K'_{0n} = \int_0^{\frac{\pi}{2}} d\phi = \frac{\pi}{2}, \quad (n = \text{limit,})$$

we have

$$(20)^* \quad (1 + k'_0)(1 + k'_{00}) \cdots (1 + k'_{0n}) = \frac{2K'}{\pi}.$$

From eq. (19)\* we have, since [eq. (10), Chap. IV]

$$\frac{1 + k}{2} = \frac{1}{1 + k'_0}$$

$$(1 + k'_0) \int_0^{\phi_0} \frac{d\phi_0}{\Delta(\phi_0, k_1)} = \int_0^{\phi} \frac{d\phi}{\Delta(\phi_1, k)};$$

whence also, since for  $\phi_0 = \frac{\pi}{2}$ ,  $\phi = \pi$ ,

$$(1 + k'_0)K_1 = 2K,$$

$$(1 + k'_{00})K_2 = 2K_1,$$

$$\dots \dots \dots$$

$$(1 + k'_{0n})K_n = 2K_{n-1},$$

and

$$(1 + k'_0)(1 + k'_{00}) \cdots (1 + k'_{0n})K_n = 2^n K;$$

or

$$(21) \quad \frac{K_n}{2^n} = \frac{K}{(1 + k'_0)(1 + k'_{00}) \cdots} \quad (n = \infty)$$

$$= \frac{\pi}{2K_1} K.$$

Let us find the limiting value of  $F(k_{0n}, \phi_n)$  in eq. (15). In the equation  $\tan(\phi_n - \phi_{n-1}) = k_{n-1} \tan \phi_{n-1}$ , we see that when  $k_{n-1}$  reaches the limit 1, then  $\phi_n - \phi_{n-1} = \phi_{n-1}$  or  $\phi_n = 2\phi_{n-1}$ . Therefore

$$\begin{aligned} \frac{\phi_n}{2^n} &= \frac{2\phi_{n-1}}{2^n} = \frac{\phi_{n-1}}{2^{n-1}}, \\ \frac{\phi_{n+1}}{2^{n+1}} &= \frac{2\phi_n}{2^{n+1}} = \frac{\phi_n}{2^n} = \frac{\phi_{n-1}}{2^{n-1}}, \\ \frac{\phi_{n+m}}{2^{n+m}} &= \frac{\phi_{n-1}}{2^{n-1}} = \text{constant, whatever } m \text{ may be.} \end{aligned}$$

Therefore eq. (15) becomes

$$(21)^* \quad F(k, \phi) = (1 + k_0)(1 + k_{00}) \cdots (1 + k_{0n}) \frac{\phi_n}{2^n},$$

$n$  being whatever number will carry  $k_0$  and  $\frac{\phi_1}{2}$  to their limiting values.

In the same way, eqs. (16) and (17) become

$$(22) \quad F(k, \phi) = \frac{1}{\cos^2 \frac{\theta}{2} \cos^2 \frac{\theta_0}{2} \cdots \cos^2 \frac{\theta_{0n}}{2}} \cdot \frac{\phi_n}{2^n}$$

$$(23) \quad = \sqrt{\frac{\cos \theta_0 \cos \theta_{00} \cdots \cos \theta_{0n}}{\cos \theta}} \cdot \frac{\phi_n}{2^n},$$

$n - 1$  being the number which makes  $k'_{n-1} = 1$ .

In these last three equations  $k_0, k_{00}$  are determined by eqs. (14<sub>1</sub>);  $\phi_1, \phi_2$ , etc., by eqs. (14<sub>2</sub>)\*;  $\theta, \theta_0$ , etc., by eqs. (14<sub>1</sub>); and  $k', k'_1, k'_2$ , etc., for use in eq. (14<sub>2</sub>) by eqs. (14<sub>1</sub>).

---

\*Taking for  $\phi_1 - \phi$ , etc., not always the least angle given by the tables, but that which is nearest to  $\phi$ .

## BISECTED AMPLITUDES.

We have identically

$$u = 2 \cdot \frac{u}{2} = 2 \int \frac{d \operatorname{am} \frac{u}{2}}{\sqrt{1 - k^2 \operatorname{sn}^2 \frac{u}{2}}};$$

$$\frac{u}{2} = 2 \cdot \frac{u}{4} = 2F \left( k, \operatorname{am} \frac{u}{4} \right);$$

etc.

Therefore

$$\begin{aligned} u &= F(k, \operatorname{am} u) = 2^n F \left( k, \operatorname{am} \frac{u}{n} \right) \\ &= 2^n \cdot \operatorname{am} \frac{u}{n}, \end{aligned} \quad (n = \text{limit,})$$

$\operatorname{am} \frac{u}{n}$  being determined by repeated applications of [eq. \(12\)](#) of Chap. II, as follows:

$$\operatorname{sn}^2 \frac{u}{2} = \frac{1 - \operatorname{cn} u}{1 + \operatorname{dn} u} = \frac{2 \sin^2 \frac{1}{2} \operatorname{am} u}{1 + \operatorname{dn} u};$$

$$(24) \quad \operatorname{sn} \frac{u}{2} = \frac{\sin \frac{1}{2} \operatorname{am} u}{\sqrt{\frac{1 + \cos \beta}{2}}} = \frac{\sin \frac{\operatorname{am} u}{2}}{\cos \frac{1}{2} \beta};$$

$\beta$  being an angle determined by the equation

$$(25) \quad \cos \beta = \operatorname{dn} u = \sqrt{1 - k^2 \operatorname{sn}^2 u},$$

and  $n$  being the number which makes

$$2^n \operatorname{am} \frac{u}{n} = \text{constant.}$$

$\text{am} \frac{u}{n}$  is found by repeated applications of eq. (24).  
 Indicating the amplitudes as follows:

$$\begin{aligned}
 & \text{am } u = \phi, \\
 & \text{am } \frac{u}{2} = \phi_{02}, \\
 & \text{am } \frac{u}{4} = \phi_{04}, \\
 (26) \quad & \text{am } \frac{u}{8} = \phi_{08}, \\
 & \dots \dots \dots \\
 & \text{am } \frac{u}{2^n} = \phi_{02^n}, \text{---} \\
 & F(k, \phi) = 2^n \phi_{02^n};
 \end{aligned}$$

$n$  being the limiting value.

In eq. (18), when  $k_n$  reaches its limit 1, we have

$$F(k_n, \phi_{0n}) = \int_0^\phi \frac{d\phi_{0n}}{\cos \phi_{0n}} = \log_\epsilon \tan(45^\circ + \frac{1}{2}\phi_{0n}),$$

and eqs. (18) and (19) become

$$\begin{aligned}
 (27) \quad F(k, \phi) &= (1 + k'_0)(1 + k'_{00}) \dots (1 + k'_{0n}) \log_\epsilon \tan(45^\circ + \frac{1}{2}\phi_{0n}) \\
 &= \frac{1}{\cos^2 \frac{1}{2}\eta \cos^2 \frac{1}{2}\eta_0 \dots \cos^2 \frac{1}{2}\eta_{0n}} \log_\epsilon \tan(45^\circ + \frac{1}{2}\phi_{0n})
 \end{aligned}$$

$$(28) \quad = \frac{1}{\cos^2 \frac{1}{2}\eta \cos^2 \frac{1}{2}\eta_0 \dots \cos^2 \frac{1}{2}\eta_{0n}} \cdot \frac{1}{M} \log \tan(45^\circ + \frac{1}{2}\phi_{0n});$$

$n$  being the number which renders  $k_n = 1$ .

Eq. (20) becomes

$$\begin{aligned}
 (29) \quad F(k, \phi) &= \sqrt{\frac{k_1 k_2 \cdots k_n^2}{k}} \cdot \log_{\epsilon} \tan(45^\circ + \frac{1}{2} \phi_{0n}) \\
 &= \sqrt{\frac{k_1 k_2 \cdots k_n^2}{k}} \cdot \frac{1}{M} \log \tan(45^\circ + \frac{1}{2} \phi_{0n}) \\
 &= \sqrt{\frac{\cos \eta_0 \cos \eta_{00} \cdots \cos^2 \eta_{0n}}{\cos \eta}} \cdot \frac{1}{M} \log \tan(45^\circ + \frac{1}{2} \phi_{0n}).
 \end{aligned}$$

In these equations  $k'_0, k'_{00}$ , etc., are determined by eqs. (18<sub>3</sub>);  $\eta, \eta_0$ , etc., by eqs. (18<sub>3</sub>);  $\phi_0, \phi_{00}$ , etc., by eqs. (18<sub>2</sub>);  $k_1, k_2$ , etc., by eqs. (18<sub>1</sub>).

Substituting in eq. (27) from eq. (20)\*, we have

$$\begin{aligned}
 (30) \quad F(k, \phi) &= \frac{2K'}{\pi} \log_{\epsilon} \tan(45^\circ + \frac{1}{2} \phi_{0n}) \\
 &= \frac{2K'}{\pi M} \log \tan(45^\circ + \frac{1}{2} \phi_{0n}).
 \end{aligned}$$



# CHAPTER V.

## COMPLETE FUNCTIONS

INDICATE by  $K$  the complete integral

$$(1) \quad K = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}},$$

and by  $K_0$  the complete integral

$$(2) \quad K_0 = \int_0^{\frac{\pi}{2}} \frac{d\phi_1}{\sqrt{1 - k_0^2 \sin^2 \phi_1}};$$

and in a similar manner  $K_{00}$ ,  $K_{03}$ , etc.

From [eq. \(12\)](#), Chap. IV, we have

$$\begin{aligned} \tan(\phi_1 - \phi) &= k' \tan \phi \\ &= \frac{\tan \phi_1 - \tan \phi}{1 + \tan \phi_1 \tan \phi} \end{aligned}$$

whence

$$\begin{aligned} \tan \phi_1 &= \frac{(1 + k') \tan \phi}{1 - k' \tan^2 \phi} \\ &= \frac{1 + k'}{\frac{1}{\tan \phi} - k' \tan \phi}. \end{aligned}$$

From this equation we see that when  $\phi = \frac{\pi}{2}$ ,  $\phi_1 = \pi$ . This same result might also have been deduced from [Fig. 1](#), Chap. IV, or from the equation

$$(3) \quad \phi_1 = 2\phi - k_0 \sin 2\phi + \frac{1}{2}k_0^2 \sin 4\phi - \text{etc.},$$

this last being the well-known trigonometrical formula

$$\tan x = n \tan y,$$

$$x = y - \frac{1-n}{1+n} \sin 2y + \frac{1}{2} \left( \frac{1-n}{1+n} \right)^2 \sin 4y - \frac{1}{3} \left( \frac{1-n}{1+n} \right)^3 \sin 6y + \text{etc.}$$

Since

$$\int_0^{\frac{\pi}{2}} \frac{d\phi_1}{\Delta(k_0, \phi_1)} = K_0, \text{ we must have}$$

$$\int_0^{\pi} \frac{d\phi_1}{\Delta(k_0, \phi_1)} = 2K_0.$$

These values substituted in eq. (13), Chap. IV, give successively

$$(4) \quad \begin{aligned} K &= (1 + k_0)K_0, \\ K_0 &= (1 + k_{00})K_{00}, \\ &\dots \dots \dots \\ K_{0(n-1)} &= (1 + k_{0n})K_{0n}; \end{aligned}$$

whence

$$(5) \quad K = (1 + k_0)(1 + k_{00}) \cdots (1 + k_{0n})K_{0n}.$$

Since the limit of  $k_{0n}$  is 0,  $K_{0n}$  becomes

$$K_{0n} = \int_0^{\frac{\pi}{2}} d\phi = \frac{\pi}{2},$$

and we have

$$(6) \quad K = \frac{\pi}{2}(1 + k_0)(1 + k_{00}) \cdots$$

$$(7) \quad = \frac{\frac{1}{2}\pi}{\cos^2 \frac{1}{2}\theta \cos^2 \frac{1}{2}\theta_0 \cdots \cos^2 \frac{1}{2}\theta_{0n}};$$

$k_1, k_0$ , etc., and  $\theta_1, \theta_0$ , etc., being found by eqs. (14<sub>1</sub>) of Chap. IV.

From the formulæ in these two chapters we can compute the values of  $u$  for all values of  $\phi$  and  $k$  and arrange them in tables. These are Legendre's Tables of Elliptic Integrals.





These methods for finding  $\phi$  apply only when  $\phi < \frac{\pi}{2}$ , that is,  $u < K$ . In the opposite case ( $u > K$ ) put

$$u = 2nK \pm v,$$

the upper or the lower sign being taken according as  $K$  is continued in  $u$  an even or an odd number of times. In either case  $v < K$ , and we can find  $v$  by the preceding methods.

Having found  $v$ , we have from eq. (5), Chap. III,

$$\begin{aligned} \operatorname{am} u &= \operatorname{am}(2nK \pm v) \\ &= n\pi \pm \operatorname{am} v. \end{aligned}$$

## CHAPTER VII.

### DEVELOPMENT OF ELLIPTIC FUNCTIONS INTO FACTORS.

From eq. (12), Chap. IV, we readily get

$$\begin{aligned}\sin(2\phi_0 - \phi) &= k \sin \phi; \\ \sin \phi &= \frac{\sin 2\phi_0}{\sqrt{1 + k^2 + 2k \cos 2\phi_0}} \\ &= \frac{\sin 2\phi_0}{\sqrt{(1 + k)^2 - 4k \sin^2 \phi_0}} \\ &= \frac{1 + k'_0}{2} \cdot \frac{\sin 2\phi_0}{\sqrt{1 - k_1^2 \sin^2 \phi_0}}\end{aligned}$$

(since  $\frac{4k}{1+k} = k_1$  and  $1+k = \frac{2}{1+k'_0}$ , eqs. (6) and (10), Chap. IV); and thence

$$(1) \quad \sin \phi = \frac{(1 + k'_0) \sin \phi_0 \cos \phi_0}{\Delta(\phi_0, k_1)}.$$

From eq. (13), Chap. IV, we have

$$\int_0^{\phi_0} \frac{d\phi_0}{\Delta(\phi_0, k_1)} = \frac{1+k}{2} \int_0^{\phi} \frac{d\phi}{\Delta(\phi, k)};$$

and from eq. (4), Chap. V, passing up the scale of moduli one step,

$$1 + k = \frac{K_1}{K},$$

whence

$$F(\phi_0, k_1) = \frac{K_1}{2K} F(\phi, k).$$

Put

$$F(\phi_0, k_1) = u_1 \quad \text{and} \quad F(\phi, k) = u,$$

whence

$$u_1 = \frac{K_1}{2K} u.$$

Furthermore,

$$\phi = \text{am}(u, k);$$

$$\phi_1 = \text{am}(u_1, k_1) = \text{am}\left(\frac{K_1}{2K} u, k_1\right).$$

Substituting these values in [eq. \(1\)](#), we have

$$\text{sn}(u, k) = (1 + k'_0) \frac{\text{sn}\left(\frac{K_1}{2K} u, k_1\right) \text{cn}\left(\frac{K_1}{2K} u, k_1\right)}{\text{dn}\left(\frac{K_1}{2K} u, k_1\right)}.$$

But from [eq. \(11\)](#), Chap. III, we have

$$\frac{\text{cn}(v, k_1)}{\text{dn}(v, k_1)} = \text{sn}(v + K_1, k_1),$$

or

$$\begin{aligned} \frac{\text{cn}\left(\frac{K_1}{2K} u, k_1\right)}{\text{dn}\left(\frac{K_1}{2K} u, k_1\right)} &= \text{sn}\left(\frac{K_1 u}{2K} + K_1, k_1\right) \\ &= \text{sn}\left(\frac{K_1}{2K} (u + 2K), k_1\right); \end{aligned}$$



whence

$$(2) \quad \operatorname{sn}(u, k) = (1 + k'_0) \operatorname{sn} \frac{K_1 u}{2K} \operatorname{sn} \left[ \frac{K_1}{2K} (u + 2K) \right].^*$$

(Mod. =  $k_1$ .)

From this equation, evidently, we have generally

$$(2)^* \quad \operatorname{sn}(v, k_n) = (1 + k'_{0(n+1)}) \operatorname{sn} \frac{K_{n+1} v}{2K_n} \operatorname{sn} \left[ \frac{K_{n+1}}{2K_n} (v + 2K_n) \right].$$

(Mod. =  $k_{n+1}$ .)

Applying this general formula to the two factors of [eq. \(2\)](#), we have

$$\begin{aligned} \operatorname{sn} \left( \frac{K_1 u}{2K}, k_1 \right) &= (1 + k'_{00}) \operatorname{sn} \frac{K_2}{2K_1} \cdot \frac{K_1 u}{2K} \cdot \operatorname{sn} \left[ \frac{K_2}{2K_1} \left( \frac{K_1 u}{2K} + 2K_1 \right) \right] \\ &= (1 + k'_{00}) \operatorname{sn} \frac{K_2 u}{2^2 K} \operatorname{sn} \frac{K_2}{2^2 K} (u + 4K); \end{aligned}$$

(Mod.  $k_2$ )  
(Mod.  $k_2$ );

$$(3) \quad \operatorname{sn} \left[ \frac{K_1}{2K} (u + 2K), k_1 \right] = (1 + k'_{00}) \operatorname{sn} \frac{K_2}{2^2 K} (u + 2K) \cdot \operatorname{sn} \frac{K_2}{2K_1} \left[ \frac{K_1}{2K} (u + 2K) + 2K_1 \right].$$

(Mod.  $k_2$ .)

---

\*The analogous formula in Trigonometry is

$$\sin \phi = 2 \sin \frac{1}{2} \phi \sin \frac{1}{2} (\phi + \pi).$$

The last argument in this equation is equal to

$$\frac{K_2}{2^2K}(u + 6K);$$

and since, [eq. \(7\)](#), Chap. III,

$$\operatorname{sn}(u, k_2) = \operatorname{sn}(2K_2 - u, k_2),$$

we can put in place of this,

$$2K_2 - \frac{K_2}{2^2K}(u + 6K) = \frac{K_2}{2^2K}(2K - u);$$

whence [eq. \(3\)](#) becomes

$$\operatorname{sn} \left[ \frac{K_1}{2K}(u + 2K), k_1 \right] = (1 + k'_{00}) \operatorname{sn} \frac{K_2}{2^2K}(2K + u) \cdot \operatorname{sn} \frac{K_2}{2^2K}(2K - u). \quad (\text{Mod. } k_2.)$$

Substituting these values in [eq. \(2\)](#), we have

$$(4) \quad \operatorname{sn}(u, k) = (1 + k'_0)(1 + k'_{00})^2 \operatorname{sn} \frac{K_2 u}{2^2K} \cdot \operatorname{sn} \frac{K_2}{2^2K}(2K \pm u) \operatorname{sn} \frac{K_2}{2^2K}(4K + u), \quad (\text{Mod. } k_2),$$

in which the double sign indicates two separate factors which are to be multiplied together.

By the application of the general [equation \(2\)\\*](#) we find that the arguments in the second member of [eq. \(4\)](#) will each give rise to two new arguments, as follows:

$$\frac{K_2 u}{2^2K} \quad \text{gives} \quad \frac{K_3 u}{2^3K'}$$

and

$$\frac{K_3}{2K_2} \left( \frac{K_2 u}{2^2 K} + 2K_2 \right) = \frac{K_3}{2^3 K} (u + 8K);$$

$$\frac{K_2}{2^2 K} (2K \pm u) \quad \text{gives} \quad \frac{K_3}{2^3 K} (2K \pm u),$$

and

$$(a) \quad \frac{K_3}{2K_2} \left[ \frac{K_2}{2^2 K} (2K \pm u) + 2K_2 \right] = \frac{K_3}{2^3 K} (10K \pm u),$$

$$\frac{K_2}{2^2 K} (4K + u) \quad \text{gives} \quad \frac{K_3}{2^3 K} (4K + u),$$

and

$$(b) \quad \frac{K_3}{2K_2} \left[ \frac{K_2}{2^2 K} (4K + u) + 2K_2 \right] = \frac{K_3}{2^3 K} (12K + u).$$

Subtracting (a) and (b) from  $2K_3$ , by which the sine of the amplitudes will not be changed [eq. (7), Chap. III], and since our new modulus is  $k_3$ , we have for the expressions (a) and (b),

$$(a') \quad \frac{K_3}{2^3 K} (6K \mp u);$$

$$(b') \quad \frac{K_3}{2^3 K} (4K - u).$$

Substituting these values in eq. (4), and remembering the factor  $(1 + k'_{03})$  introduced by each application of eq. (2)\*, we have

$$\begin{aligned} \text{sn}(u, k) &= (1 + k'_0)(1 + k'_{00})^2(1 + k'_{03})^4 \text{sn} \frac{K_3 u}{2^3 K} \\ &\cdot \text{sn} \frac{K_3}{2^3 K} (2K \pm u) \text{sn} \frac{K_3}{2^3 K} (4K \pm u) \\ &\cdot \text{sn} \frac{K_3}{2^3 K} (6K \pm u) \text{sn} \frac{K_3}{2^3 K} (8K + u). \end{aligned} \quad (\text{Mod. } k_3.)$$

From this the law governing the arguments is clear, and we can write for the general equation

$$\begin{aligned}
 (5) \quad \operatorname{sn}(u, k) &= (1 + k'_0)(1 + k'_{00})^2(1 + k'_{03})^4 \cdots (1 + k'_{0n})^{2^{n-1}} \\
 &\cdot \operatorname{sn} \frac{K_n u}{2^n K} \operatorname{sn} \frac{K_n}{2^n K} (2K \pm u) \\
 &\cdot \operatorname{sn} \frac{K_n}{2^n K} (4K \pm u) \operatorname{sn} \frac{K_n}{2^n K} (6K \pm u) \\
 &\cdots \operatorname{sn} \frac{K_n}{2^n K} \left[ (2^n - 2)K \pm u \right] \\
 &\cdot \operatorname{sn} \frac{K_n}{2^n K} (2^n K + u). \qquad (\text{Mod. } k_n.)
 \end{aligned}$$

Indicate the continued product of the binomial factors by  $A'$ , and we have

$$A' = (1 + k'_0)(1 + k'_{00})^2(1 + k'_{03})^4(1 + k'_{04})^8 \cdots$$

Since the limit of  $k'_0, k'_{00},$  etc., is zero, it is evident that these factors converge toward the value unity. It can be shown that the functional factors also converge toward the value unity. Thus the argument of the last factor can be written

$$K_n + \frac{K_n u}{2^n K}.$$

From [eq. \(11\)](#), Chap. III, we get then

$$\operatorname{sn} \left( K_n + \frac{K_n u}{2^n K} \right) = \frac{\operatorname{cn} \frac{K_n u}{2^n K}}{\operatorname{dn} \frac{K_n u}{2^n K}}. \qquad (\text{Mod. } k_n.)$$

But since  $k_n$  at its limit is equal to unity,  $\operatorname{cn} = \operatorname{dn}$ ; whence the last factor of [eq. \(5\)](#) is unity.

From [eq. \(21\)](#), Chap. IV, we have

$$\operatorname{limit} \frac{K_n}{2^n K} = \frac{2\pi}{2K'}.$$

Therefore for  $n = \infty$ , eq. (5) becomes

$$\begin{aligned} \operatorname{sn}(u, k) = A' \operatorname{sn} \frac{\pi u}{2K'} \operatorname{sn} \frac{\pi}{2K'} (2K \pm u) \\ \cdot \operatorname{sn} \frac{\pi}{2K'} (4K \pm u) \operatorname{sn} \frac{\pi}{2K'} (6K \pm u), \dots \end{aligned} \quad (\text{Mod. } 1),$$

or

$$(6) \quad \operatorname{sn}(u, k) = A' \operatorname{sn} \frac{\pi u}{2K'} \left[ \prod_1^{\infty} h \right] \operatorname{sn} \frac{\pi}{2K'} (2hK \pm u), \quad (\text{Mod. } 1),$$

where the sign  $[\prod]$  indicates the continued product in the same manner as  $\Sigma$  indicates the continued sum.

When  $k = 1$ ,  $\int_0^\phi F(\phi, k) d\phi$  becomes

$$v = \int_0^\phi \frac{d\phi}{\cos \phi} = \frac{1}{2} \log_{\epsilon} \frac{1 + \sin \phi}{1 - \sin \phi};$$

whence

$$e^{2v} = \frac{1 + \sin \phi}{1 - \sin \phi},$$

and

$$\sin \phi = \frac{e^{2v} - 1}{e^{2v} + 1} = \frac{e^v - e^{-v}}{e^v + e^{-v}} = \operatorname{sn}(v, 1).$$

Hence in equation (6)

$$\begin{aligned} \sin \frac{\pi u}{2K'} &= \frac{e^{\frac{\pi u}{2K'}} - e^{-\frac{\pi u}{2K'}}}{e^{\frac{\pi u}{2K'}} + e^{-\frac{\pi u}{2K'}}}; \\ \operatorname{sn} \frac{\pi}{2K'} (2hK \pm u) &= \frac{e^{\frac{h\pi K}{K'}} e^{\pm \frac{\pi u}{2K'}} - e^{-\frac{h\pi K}{K'}} e^{\mp \frac{\pi u}{2K'}}}{e^{\frac{h\pi K}{K'}} e^{\pm \frac{\pi u}{2K'}} + e^{-\frac{h\pi K}{K'}} e^{\mp \frac{\pi u}{2K'}}}. \end{aligned}$$

Put

$$(6)^* \quad q = e^{-\frac{\pi K'}{K}}, \quad q' = e^{-\frac{\pi K}{K'}},$$

and the last expression becomes

$$\begin{aligned} \operatorname{sn} \frac{\pi}{2K'}(2hK \pm u) &= \frac{q'^{-h} e^{\pm \frac{\pi u}{2K'}} - q'^h e^{\mp \frac{\pi u}{2K'}}}{q'^{-h} e^{\pm \frac{\pi u}{2K'}} + q'^h e^{\mp \frac{\pi u}{2K'}}}; \\ \operatorname{sn} \frac{\pi}{2K'}(2hK + u) \operatorname{sn} \frac{\pi}{2K'}(2hK - u) &= \frac{q'^{-h} e^{\frac{\pi u}{2K'}} - q'^h e^{-\frac{\pi u}{2K'}}}{q'^{-h} e^{\frac{\pi u}{2K'}} + q'^h e^{-\frac{\pi u}{2K'}}} \cdot \frac{q'^{-h} e^{-\frac{\pi u}{2K'}} - q'^h e^{\frac{\pi u}{2K'}}}{q'^{-h} e^{-\frac{\pi u}{2K'}} + q'^h e^{\frac{\pi u}{2K'}}} \\ &= \frac{q'^{-2h} + q'^{2h} - \left( e^{\frac{\pi u}{K'}} + e^{-\frac{\pi u}{K'}} \right)}{q'^{-2h} + q'^{2h} + \left( e^{\frac{\pi u}{K'}} + e^{-\frac{\pi u}{K'}} \right)}. \end{aligned}$$

From plane trigonometry we have the equations

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = -i \tan ix, \quad e^x + e^{-x} = 2 \cos ix;$$

where  $i = \sqrt{-1}$ : which gives

$$\operatorname{sn} \frac{\pi u}{2K'} = -i \tan \frac{\pi i u}{2K'}; \quad (\text{Mod. } 1;)$$

$$\operatorname{sn} \frac{\pi}{2K'}(2hK + u) \operatorname{sn} \frac{\pi}{2K'}(2hK - u)$$

$$= \frac{q'^{-2h} + q'^{2h} - 2 \cos \frac{\pi i u}{K'}}{q'^{-2h} + q'^{2h} + 2 \cos \frac{\pi i u}{K'}}$$

$$(7) \quad = \frac{1 - 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}{1 + 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}.$$

From [eq. \(10\)](#), Chap. III, we have

$$\operatorname{sn}(u, k) = -i \operatorname{tn}(iu, k').$$

Substituting these values in [eq. \(6\)](#), we have

$$\operatorname{tn}(iu, k') = A' \tan \frac{\pi i u}{2K'} [\Pi] \frac{1 - 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}{1 + 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}.$$

Now in place of the series of moduli  $k'$ ,  $k'_0$  and the corresponding complete integral  $K'$ , we are at liberty to substitute the parallel series of moduli  $k$ ,  $k_0$  and the corresponding complete integral  $K$ ; calling the

new integral  $u$ , we have

$$\begin{aligned}
 (8) \quad \operatorname{tn}(u, k) &= A \tan \frac{\pi u}{2K} [\Pi] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}} \\
 &= A \tan \frac{\pi u}{2K} \frac{1 - 2q^2 \cos \frac{\pi u}{K} + q^4}{1 + 2q^2 \cos \frac{\pi u}{K} + q^4} \\
 &\quad \cdot \frac{1 - 2q^4 \cos \frac{\pi u}{K} + q^8}{1 + 2q^4 \cos \frac{\pi u}{K} + q^8} \\
 &\quad \cdot \frac{1 - 2q^6 \cos \frac{\pi u}{K} + q^{12}}{1 + 2q^6 \cos \frac{\pi u}{K} + q^{12}} \cdots,
 \end{aligned}$$

where

$$(9) \quad A = (1 + k_0)(1 + k_{00})^2(1 + k_{03})^4(1 + k_{04})^8 \dots$$

Now in [equation \(6\)](#) put  $u + K$  for  $u$ , and we have, since [[eq. \(11\)](#), Chap. III]  $\operatorname{sn}(u + K) = \frac{\operatorname{cn} u}{\operatorname{dn} u'}$ ,

$$\begin{aligned}
 \frac{\operatorname{cn} u}{\operatorname{dn} u} &= A' \operatorname{sn} \frac{\pi(u + K)}{2K'} \\
 &[\Pi] \operatorname{sn} \frac{\pi}{2K'} [(2h + 1)K + u] \cdot \operatorname{sn} \frac{\pi}{2K'} [(2h - 1)K - u].
 \end{aligned}$$

(Mod. 1.)

Now from  $2h - 1$  and  $2h + 1$  we have the following series of numbers respectively:

$$\begin{aligned}
 2h - 1 : & \quad 1, \quad 3, \quad 5, \quad 7, \quad 9, \quad \text{etc.} \\
 2h + 1 : & \quad 3, \quad 5, \quad 7, \quad 9, \quad \text{etc.}
 \end{aligned}$$



It will be observed that the factor outside of the sign  $[\Pi]$ , viz.,  $\sin \operatorname{am} \frac{\pi(u+K)}{2K'}$ , would, if placed under the sign  $[\Pi]$ , supply the missing first term of the second series. Hence, placing this factor within the sign, we have

$$(10) \quad \frac{\operatorname{cn} u}{\operatorname{dn} u} = A' [\Pi] \operatorname{sn} \frac{\pi}{2K'} [(2h-1)K+u] \cdot \operatorname{sn} \frac{\pi}{2K'} [(2h-1)K-u].$$

(Mod. 1.)

Comparing this with [equation \(7\)](#), we see that the factors herein differ from those in [equation \(7\)](#) only in having  $2h-1$  in place of  $2h$ ; hence we have

$$\begin{aligned} & \operatorname{sn} \frac{\pi}{2K'} [(2h-1)K+u] \operatorname{sn} \frac{\pi}{2K'} [(2h-1)K-u] \\ &= \frac{1 - 2q^{2h-1} \cos \frac{\pi i u}{K'} + q^{4h-2}}{1 + 2q^{2h-1} \cos \frac{\pi i u}{K'} + q^{4h-2}}. \end{aligned}$$

(Mod. 1.)

From [eqs. \(10\)](#), Chap. III, we have

$$\frac{\operatorname{cn}(u, k)}{\operatorname{dn}(u, k)} = \frac{1}{\operatorname{dn}(iu, k')};$$

whence [eq. \(10\)](#) becomes

$$(11) \quad \frac{1}{\operatorname{dn}(iu, k')} = A' [\Pi] \frac{1 - 2q^{2h-1} \cos \frac{\pi i u}{K'} + q^{4h-2}}{1 + 2q^{2h-1} \cos \frac{\pi i u}{K'} + q^{4h-2}};$$

and when in place of  $iu, k', K', q', A'$ , we substitute  $u, k, K, q$  and  $A$ , and invert the equation, we have

$$(12) \quad \operatorname{dn}(u, k) = \frac{1}{A} [\Pi] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

Bearing in mind the remarkable property (Chap. III, p. 31) that the functions  $\operatorname{sn} u$  and  $\operatorname{dn} u$  approach infinity for the same value of  $u$ , we see that both these functions, except as to the factor independent of  $u$ , must have the same denominator. Furthermore, since  $\operatorname{sn} u$  and  $\operatorname{tn} u$  disappear for the same value of  $u$ , they must, except for the independent factor, have the same numerator. Hence, indicating by  $B$  a new quantity, dependent upon  $k$  but independent of  $u$ , we have

$$(13) \quad \operatorname{sn}(u, k) = B \sin \frac{\pi u}{2K} [\Pi] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

and since

$$\operatorname{cn} u = \frac{\operatorname{sn} u}{\operatorname{tn} u},$$

we also have, from eqs. (8) and (13),

$$(14) \quad \operatorname{cn}(u, k) = \frac{B}{A} \cos \frac{\pi u}{2K} [\Pi] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

Collecting these results, we have the following equations:

$$(15) \quad \operatorname{sn}(u, k) = B \sin \frac{\pi u}{2K} [\Pi] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}},$$

$$(16) \quad \operatorname{cn}(u, k) = \frac{B}{A} \cos \frac{\pi u}{2K} [\Pi] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}},$$

$$(17) \quad \operatorname{dn}(u, k) = \frac{1}{A} [\Pi] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

To ascertain the values of  $A$  and  $B$ , we proceed as follows:

In eq. (17) we make  $u = 0$ , whence, by eq. (13), Chap. II, we have

$$1 = \frac{1}{A} [\Pi] \left( \frac{1 + q^{2h-1}}{1 - q^{2h-1}} \right)^2;$$

whence

$$(18) \quad \frac{1}{A} = [\Pi] \left( \frac{1 - q^{2h-1}}{1 + q^{2h-1}} \right)^2.$$

In equation (17), making  $u = K$ , we get, by equation (1), Chap. III,

$$(19) \quad k' = \frac{1}{A} [\Pi] \left( \frac{1 - q^{2h-1}}{1 + q^{2h-1}} \right)^2 = \frac{1}{A^2};$$

$$\therefore \frac{1}{A} = \sqrt{k'}.$$

We have identically

$$1 = B \frac{1}{B} = B \frac{\frac{A}{B}}{\frac{A}{A}} = B \frac{\sqrt{k'}}{\frac{A}{A}};$$

whence

$$\frac{B}{A} = B\sqrt{k'}.$$

To calculate  $B$ , put  $e^{\frac{i\pi u}{2K}} = v$ ; if we change  $\frac{\pi u}{2K}$  into  $\frac{\pi u}{2K} + \frac{i\pi K'}{2K}$ ,  $v$  will change into  $v\sqrt{q}$ , and  $\text{sn } u$  will become, by eq. (14), Chap. III,

$$\text{sn}(u + iK') = \frac{1}{k \text{sn } u}.$$

Now, replacing  $\sin \frac{\pi u}{2K}$  and  $\cos \frac{\pi u}{K}$  by their exponential values, and observing that

$$1 - 2q^n \cos \frac{\pi u}{K} + q^{2n} = (1 - q^n v^2)(1 - q^n v^{-2}),$$

we have

$$\operatorname{sn} u = \frac{B}{2} \cdot \frac{v - v^{-1}}{\sqrt{-1}} \cdot \frac{[\Pi](1 - q^{2h} v^2)(1 - q^{2h} v^{-2})}{[\Pi](1 - q^{2h-1} v^2)(1 - q^{2h-1} v^{-2})}.$$

Changing  $u$  into  $u + iK'$ , and consequently  $v$  into  $v\sqrt{q}$ , we have

$$\frac{1}{k \operatorname{sn} u} = \frac{B}{2} \cdot \frac{v\sqrt{q} - v^{-1}\sqrt{q^{-1}}}{\sqrt{-1}} \cdot \frac{[\Pi](1 - q^{2h+1} v^2)(1 - q^{2h-1} v^{-2})}{[\Pi](1 - q^{2h} v^2)(1 - q^{2h-2} v^{-2})}.$$

Multiplying these equations together, member by member, and observing that

$$\begin{aligned} v\sqrt{q} - v^{-1}\sqrt{q^{-1}} &= \frac{1 - qv^2}{-v\sqrt{q}}, \\ v - v^{-1} &= v(1 - v^{-2}), \end{aligned}$$

we get

$$\begin{aligned} \frac{1}{k} &= \frac{B^2}{4} \cdot \frac{1 - qv^2}{v\sqrt{q}} \cdot v(1 - v^{-2}) \cdot \frac{[\Pi](1 - q^{2h+1} v^2)(1 - q^{2h} v^{-2})}{[\Pi](1 - q^{2h-1} v^2)(1 - q^{2h-2} v^{-2})} \\ &= \frac{B^2}{4\sqrt{q}} (1 - qv^2)v(1 - v^{-2}) \frac{(1 - q^3 v^2)(1 - q^5 v^2) \dots}{(1 - qv^2)(1 - q^3 v^2) \dots} \\ &\quad \cdot \frac{(1 - q^2 v^{-2})(1 - q^4 v^{-2}) \dots}{(1 - v^{-2})(1 - q^2 v^{-2}) \dots} \\ &= \frac{B^2}{4} \cdot \frac{1}{\sqrt{q}}. \end{aligned}$$

$$\therefore B = \frac{2\sqrt[4]{q}}{\sqrt{k}};$$

whence

$$\frac{B}{A} = 2\sqrt[4]{q}\sqrt{\frac{k'}{k}}.$$

Substituting these values in eqs. (15), (16), and (17), we have

$$(20) \quad \operatorname{sn}(u, k) = \frac{2\sqrt[4]{q}}{\sqrt{k}} \sin \frac{\pi u}{2K} [\Pi] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

$$(21) \quad \operatorname{cn}(u, k) = \frac{2\sqrt{k'}\sqrt[4]{q}}{\sqrt{k}} \cos \frac{\pi u}{2K} [\Pi] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

$$(22) \quad \operatorname{dn}(u, k) = \sqrt{k'} [\Pi] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

# CHAPTER VIII.

## THE $\Theta$ FUNCTION.

WE will indicate the denominator in eq. (20), Chap. VII, by  $\phi(u)$ , thus:

$$(1) \quad \phi(u) = [\Pi](1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}).$$

We will now develop this into a series consisting of the cosines of the multiples of  $\frac{\pi u}{K}$ . Put  $\frac{\pi u}{2K} = x$ , whence

$$2 \cos \frac{\pi u}{K} = (e^{2ix} + e^{-2ix});$$

but

$$1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2} = (1 - q^{2h-1} e^{2ix})(1 - q^{2h-1} e^{-2ix}),$$

and therefore

$$(2) \quad \phi(u) = (1 - qe^{2ix})(1 - q^3e^{2ix})(1 - q^5e^{2ix}) \dots \\ (1 - qe^{-2ix})(1 - q^3e^{-2ix})(1 - q^5e^{-2ix}) \dots$$

Putting now  $u + 2iK'$  instead of  $u$ , we have

$$x_1 = \frac{\pi(u + 2iK')}{2K} = x + \frac{\pi iK'}{K}, \\ 2ix_1 = 2ix - \frac{2\pi K'}{K};$$

and

$$e^{2ix_1} = q^2 e^{2ix}, \\ e^{-2ix_1} = \frac{1}{q^2} e^{-2ix}.$$

From these we have

$$\begin{aligned} \phi(u + 2iK') &= -\frac{1}{q}e^{-2ix}(1 - qe^{2ix})(1 - q^3e^{2ix}) \dots \\ &\quad (1 - qe^{-2ix})(1 - q^3e^{-2ix}) \dots ; \end{aligned}$$

whence

$$\phi(u + 2iK') = -\frac{1}{q}e^{-2ix}\phi(u),$$

or

$$(3) \quad \phi(u + 2iK') = -q^{-1}e^{-\frac{\pi iu}{K}}\phi(u).$$

Now put

$$(4) \quad \phi(u) = A + B \cos \frac{\pi u}{K} + C \cos \frac{2\pi u}{K} + D \cos \frac{3\pi u}{K} + \text{etc.}$$

Since

$$\cos \frac{\pi u}{K} = \frac{1}{2} \left( e^{2ix} + e^{-2ix} \right),$$

this becomes

$$(5) \quad \begin{aligned} \phi(u) &= A + \frac{1}{2}Be^{2ix} + \frac{1}{2}Ce^{4ix} + \frac{1}{2}De^{6ix} + \dots \\ &\quad + \frac{1}{2}Be^{-2ix} + \frac{1}{2}Ce^{-4ix} + \frac{1}{2}De^{-6ix} + \dots ; \end{aligned}$$

whence

$$(6) \quad \begin{aligned} -\frac{1}{q}e^{-2ix}\phi(u) &= -\frac{A}{q}e^{-2ix} - \frac{B}{2q} - \frac{C}{2q}e^{2ix} - \frac{D}{2q}e^{4ix} - \dots \\ &\quad - \frac{B}{2q}e^{-4ix} - \frac{C}{2q}e^{-6ix} - \frac{D}{2q}e^{-8ix} - \dots. \end{aligned}$$

Now in [equation \(5\)](#) put  $u + 2iK'$  in place of  $u$ , remembering that  $e^{2ix}$  and  $e^{-2ix}$  are thereby changed respectively into  $q^2e^{2ix}$  and  $q^{-2}e^{-2ix}$ , and we have

$$(7) \quad \begin{aligned} \phi(u + 2iK') = & A + \frac{Bq^2}{2}e^{2ix} + \frac{Cq^4}{2}e^{4ix} + \frac{Dq^6}{2}e^{6ix} + \dots \\ & + \frac{B}{2q^2}e^{-2ix} + \frac{C}{2q^4}e^{-4ix} + \dots . \end{aligned}$$

Since [equations \(6\)](#) and [\(7\)](#) are equal, we have

$$\begin{aligned} -\frac{B}{2q} &= A, & B &= -2qA; \\ -\frac{C}{2q} &= \frac{Bq^2}{2}, & C &= +2q^4A; \\ -\frac{D}{2q} &= \frac{Cq^4}{2}, & D &= -2q^9A; \\ \dots & & \dots & \end{aligned}$$

whence

$$(8) \quad \left\{ \begin{aligned} & [\Pi] (1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}) \\ & = A(1 - 2q \cos \frac{\pi u}{K} + 2q^4 \cos \frac{2\pi u}{K} - 2q^9 \cos \frac{3\pi u}{K} \\ & \quad + 2q^{16} \cos \frac{4\pi u}{K} - \dots). \end{aligned} \right.$$

The series in the second member has been designated by Jacobi and subsequent writers by  $\Theta(u)$ , thus:

$$(9) \quad \Theta(u) = 1 - 2q \cos \frac{\pi u}{K} + 2q^4 \cos \frac{2\pi u}{K} - \dots$$



# CHAPTER IX.

## THE $\Theta$ AND $H$ FUNCTIONS.

IN [equation \(20\)](#), Chap. VII, viz.,

$$\operatorname{sn}(u, k) = \frac{2\sqrt[4]{q}}{\sqrt{k}} \sin \frac{\pi u}{2K} [\Pi] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}},$$

the numerator and the denominator have been considered separately by Jacobi, who gave them a special notation and developed from them a theory second only in importance to the elliptic functions themselves.

Put [see [equation \(8\)](#), Chap. VIII]

$$(1) \quad \Theta(u) = \frac{1}{A} [\Pi] (1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}).$$

$$(2) \quad H(u) = 2 \frac{1}{A} \sqrt[4]{q} \sin \frac{\pi u}{2K} [\Pi] (1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h});$$

$A$  being a constant whose value is to be determined later. From these we have

$$(3) \quad \operatorname{sn}(u, k) = \frac{1}{\sqrt{k}} \cdot \frac{H(u)}{\Theta(u)}.$$

The functions  $\operatorname{sn} u$  and  $\operatorname{cn} u$  can also be expressed in terms of the new functions; thus we have

$$(4) \quad \operatorname{cn}(u, k) = \sqrt{\frac{k'}{k}} \cdot 2\sqrt[4]{q} \cos \frac{\pi u}{2K} [\Pi] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

or, since  $\sin x = \cos\left(x + \frac{\pi}{2}\right)$  and  $\cos x = -\sin\left(x + \frac{\pi}{2}\right)$ , and putting  $u = \frac{2Kx}{\pi}$ ,

$$\begin{aligned} \operatorname{cn}\left(\frac{2Kx}{\pi}, k\right) &= \sqrt{\frac{k'}{k}} \frac{H\left[\frac{2K}{\pi}\left(x + \frac{\pi}{2}\right)\right]}{\Theta\left(\frac{2Kx}{\pi}\right)} \\ &= \sqrt{\frac{k'}{k}} \frac{H\left[\frac{2Kx}{\pi} + K\right]}{\Theta\left(\frac{2Kx}{\pi}\right)}. \end{aligned}$$

Replacing  $\frac{2Kx}{\pi}$  by its value,  $u$ , we have

$$(5) \quad \operatorname{cn}(u, k) = \sqrt{\frac{k'}{k}} \frac{H(u + K)}{\Theta(u)}.$$

Furthermore,

$$(6) \quad \operatorname{dn}(u, k) = \sqrt{k'} [\Pi] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}$$

gives in the same manner

$$\operatorname{dn} \frac{2Kx}{\pi} = \sqrt{k'} \frac{\Theta\left[\frac{2K}{\pi}\left(x + \frac{\pi}{2}\right)\right]}{\Theta\left(\frac{2Kx}{\pi}\right)},$$

or

$$(7) \quad \operatorname{dn}(u, k) = \sqrt{k'} \frac{\Theta(u + K)}{\Theta(u)}.$$

If we put

$$(8) \quad H(u + K) = H_1(u),$$

$$(9) \quad \Theta(u + K) = \Theta_1(u),$$

the three elliptic functions can be expressed by the following formulas:

$$(10) \quad \operatorname{sn}(u, k) = \frac{1}{\sqrt{k}} \cdot \frac{H(u)}{\Theta(u)};$$

$$(11) \quad \operatorname{cn}(u, k) = \sqrt{\frac{k'}{k}} \cdot \frac{H_1(u)}{\Theta(u)};$$

$$(12) \quad \operatorname{dn}(u, k) = \sqrt{k'} \frac{\Theta_1(u)}{\Theta(u)}.$$

These functions  $\Theta$  and  $H$  can be expressed in terms of each other. By definition,

$$H(u) = 2C \sqrt[4]{q} \sin \frac{\pi u}{2K} [\Pi] \left( 1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h} \right);$$

but

$$\begin{aligned} 1 - 2q^h \cos \frac{\pi u}{K} + q^{2h} &= \left( 1 - q^h e^{\frac{\pi u \sqrt{-1}}{K}} \right) \left( 1 - q^h e^{-\frac{\pi u \sqrt{-1}}{K}} \right) \\ \sin \frac{\pi u}{2K} &= \frac{e^{\frac{\pi i u}{2K}} - e^{-\frac{\pi i u}{2K}}}{2\sqrt{-1}} \\ &= e^{-\frac{\pi i u}{2K}} \frac{1 - e^{\frac{\pi i u}{K}}}{2} \sqrt{-1}, \end{aligned}$$

and consequently

$$(13) \quad H(u) = C \sqrt[4]{q} e^{-\frac{\pi i u}{2K}} \sqrt{-1} \left( 1 - e^{\frac{\pi i u}{K}} \right) \left( 1 - q^2 e^{-\frac{\pi i u}{K}} \right) \left( 1 - q^2 e^{\frac{\pi i u}{K}} \right) \dots$$

Now, changing  $u$  into  $u + iK'$ , and remembering that  $e^{-\frac{\pi K'}{K}} = q$ , we have

$$(14) \quad H(u + iK') = Cq^{-\frac{1}{4}}e^{-\frac{\pi iu}{2K}}\sqrt{-1}\left(1 - qe^{\frac{\pi iu}{K}}\right)\left(1 - qe^{-\frac{\pi iu}{K}}\right) \\ \left(1 - q^3e^{\frac{\pi iu}{K}}\right)\left(1 - q^3e^{-\frac{\pi iu}{K}}\right)\cdots;$$

and reuniting the factors two by two, this becomes

$$(15) \quad H(u + iK') = C\sqrt{-1}q^{-\frac{1}{4}}e^{-\frac{\pi iu}{2K}} \\ \left(1 - 2q\cos\frac{\pi u}{K} + q^2\right)\left(1 - 2q^3\cos\frac{\pi u}{K} + q^6\right)\cdots;$$

and finally, according to [equation \(1\)](#),

$$(16) \quad H(u + iK') = \sqrt{-1}q^{-\frac{1}{4}}e^{-\frac{\pi iu}{2K}}\Theta(u).$$

In the same manner, we can get

$$(17) \quad \Theta(u + iK') = \sqrt{-1}q^{-\frac{1}{4}}e^{-\frac{\pi iu}{2K}}H(u).$$

Substituting  $u + 2K$  for  $u$  in [equations \(1\)](#) and [\(2\)](#), we get

$$(18) \quad \Theta(u + 2K) = \Theta(u),$$

$$(19) \quad H(u + 2K) = -H(u),$$

since  $\cos\frac{\pi}{K}(u + 2K) = \cos\frac{\pi u}{K}$  and  $\sin\frac{\pi}{2K}(u + 2K) = -\sin\frac{\pi u}{2K}$ .

The comparison of these four equations with [equations \(10\)](#), [\(11\)](#), and [\(12\)](#) shows the periodicity of the elliptic functions. For example, comparing [eqs. \(10\)](#) and [\(16\)](#) and [\(17\)](#), we see that changing  $u$  into  $u + iK'$  simply multiplies the numerator and denominator of the second

member of eq. (10) by the same number, and does not change their ratio.

The addition of  $2K$  changes the sign of the function, but not its value.

We will define  $\Theta_1$  and  $H_1$  as follows:

$$(20) \quad \Theta_1(x) = \Theta(x + K);$$

$$(21) \quad H_1(x) = H(x + K).$$

Hence we get, from equation (17),

$$\begin{aligned} \Theta_1(x + iK') &= \Theta(x + iK' + K) = \Theta(x + K + iK') \\ &= iH(x + K)e^{-\frac{i\pi}{4K}(2x+2K+iK')} \\ &= iH_1(x)e^{-\frac{i\pi}{4K}(2x+iK')}(-\sqrt{-1}), \end{aligned}$$

since 
$$e^{-\frac{i\pi}{2}} = \cos \frac{\pi}{2} - \sqrt{-1} \sin \frac{\pi}{2} = -\sqrt{-1};$$

whence

$$(22) \quad \Theta_1(x + iK') = H_1(x)e^{-\frac{i\pi}{4K}(2x+iK')}.$$

In a similar manner we get

$$(22)^* \quad H_1(x + iK') = \Theta_1(x)e^{-\frac{i\pi}{4K}(2x+iK')}.$$

In eq. (9), Chap. VIII, put  $u = \frac{2Kz}{\pi}$ , and we get

$$(23) \quad \Theta\left(\frac{2Kz}{\pi}\right) = 1 - 2q \cos 2z + 2q^4 \cos 4z - \dots .$$

Now, in this equation, changing  $z$  into  $z + \frac{\pi}{2}$ , and observing eq. (20), we get

$$(24) \quad \Theta_1\left(\frac{2Kz}{\pi}\right) = 1 + 2q \cos 2z + 2q^4 \cos 4z + \dots .$$

Applying eq. (22) to this, we have

$$\begin{aligned}
H_1\left(\frac{2Kz}{\pi}\right) &= \Theta_1\left(\frac{2K}{\pi}\left(z + \frac{\pi iK'}{2K}\right)\right) e^{\frac{\pi i}{4K}\left(\frac{4Kz}{\pi} + iK'\right)} \\
&= \Theta_1\left(\frac{2K}{\pi}\left(z + \frac{\pi iK'}{2K}\right)\right) e^{iz} q^{\frac{1}{4}} \\
&= e^{iz} q^{\frac{1}{4}} \left[ 1 + 2q \cos 2\left(z + \frac{\pi iK'}{2K}\right) + 2q^4 \cos 4\left(z + \frac{\pi iK'}{2K}\right) + \dots \right] \\
&= e^{iz} q^{\frac{1}{4}} \left[ 1 + q\left(e^{2i\left(z + \frac{\pi iK'}{2K}\right)} + e^{-2i\left(z + \frac{\pi iK'}{2K}\right)}\right) \right. \\
&\quad \left. + q^4\left(e^{4i\left(z + \frac{\pi iK'}{2K}\right)} + e^{-4i\left(z + \frac{\pi iK'}{2K}\right)}\right) + \dots \right] \\
&= e^{iz} q^{\frac{1}{4}} \left[ 1 + q(qe^{2iz} + q^{-1}e^{-2iz}) + q^4(q^2e^{4iz} + q^{-2}e^{-4iz}) + \dots \right] \\
&= e^{iz} q^{\frac{1}{4}} \left[ 1 + q^2e^{2iz} + q^6e^{4iz} + \dots + e^{-2iz} + q^2e^{-4iz} + \dots \right] \\
&= q^{\frac{1}{4}} \left[ e^{iz} + q^2e^{3iz} + q^6e^{5iz} + \dots + e^{-iz} + q^2e^{-3iz} + q^6e^{-5iz} + \dots \right] \\
&= 2q^{\frac{1}{4}} \left[ \cos z + q^2 \cos 3z + q^6 \cos 5z + \dots \right];
\end{aligned}$$

whence

$$(25) \quad H_1\left(\frac{2Kz}{\pi}\right) = 2\sqrt[4]{q} \cos z + 2\sqrt[4]{q^9} \cos 3z + 2\sqrt[4]{q^{25}} \cos 5z + \dots$$

In this equation, changing  $z$  into  $z - \frac{\pi}{2}$ , and applying eq. (21), we get

$$(26) \quad H\left(\frac{2Kz}{\pi}\right) = 2\sqrt[4]{q} \sin z - 2\sqrt[4]{q^9} \sin 3z + 2\sqrt[4]{q^{25}} \sin 5z - \dots,$$

since

$$H_1\left(\frac{2Kz}{\pi}\right) = H\left(\frac{2Kz}{\pi} + K\right).$$

We will now determine the constant  $A$  of eq. (8), Chap. VIII, and eqs. (1) and (2) of this chapter. Denote  $A$  by  $f(q)$ , and we have

$$(26)^* \quad [\Pi] (1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}) = f(q)\Theta(u).$$

Substituting herein  $u = 0$  and  $u = \frac{K}{2}$ , we have

$$[\Pi] (1 - q^{2h-1})^2 = f(q)\Theta(0);$$

$$[\Pi] (1 + q^{4h-2}) = f(q)\Theta\left(\frac{K}{2}\right).$$

From eq. (9), Chap. VIII, we get

$$(27) \quad \Theta(0) = 1 - 2q + 2q^4 - 2q^9 + 2q^{16} - \dots;$$

$$(28) \quad \Theta\left(\frac{K}{2}\right) = 1 - 2q^4 + 2q^{16} - 2q^{36} + 2q^{64} - \dots;$$

from which we see that  $\Theta(0)$  is changed into  $\Theta\left(\frac{K}{2}\right)$  when we put  $q^4$  in place of  $q$ .

Whence

$$[\Pi] (1 - q^{8h-4})^2 = f(q^4)\Theta\left(\frac{K}{2}\right);$$

and therefore

$$(29) \quad \begin{aligned} \frac{f(q)}{f(q^4)} &= [\Pi] \frac{1 + q^{4h-2}}{(1 - q^{8h-4})^2} \\ &= [\Pi] \frac{1}{(1 - q^{8h-4})(1 - q^{4h-2})}. \end{aligned}$$

Now, the expressions  $4h - 2$ ,  $8h - 4$ , and  $8h$  give the following series of numbers:

$4h - 2,$      $2,$      $6,$      $10,$      $14,$      $18,$      $22,$      $26,$      $30,$      $34;$   
 $8h - 4,$          $4,$                  $12,$                  $20,$                  $28,$                  $36;$   
 $8h,$                  $8,$                  $16,$                  $24,$                  $32.$

Hence, the three expressions taken together contain all the even numbers, and

$$[\Pi](1 - q^{8h-4})(1 - q^{4h-2})(1 - q^{8h}) = [\Pi](1 - q^{2h}).$$

Therefore, multiplying eq. (29) by

$$[\Pi] \frac{1 - q^{8h}}{1 - q^{8h}},$$

we have

$$\frac{f(q)}{f(q^4)} = [\Pi] \frac{1 - q^{8h}}{1 - q^{2h}}.$$

Now in this equation, by successive substitutions of  $q^4$  for  $q$ , we get

$$\begin{aligned} \frac{f(q^4)}{f(q^{16})} &= [\Pi] \frac{1 - q^{32h}}{1 - q^{8h}}; \\ \frac{f(q^{16})}{f(q^{64})} &= [\Pi] \frac{1 - q^{128h}}{1 - q^{32h}}; \\ \frac{f(q^{64})}{f(q^{256})} &= [\Pi] \frac{1 - q^{512h}}{1 - q^{128h}}; \\ &\dots \end{aligned}$$

Now  $q$  being less than 1,  $q^n$  tends towards the limit 0 as  $n$  increases, and consequently  $1 - q^n$  tends towards the limit 1. Also, from eq. (8),



Chap. VIII, we see that  $f(0) = 1$ . Hence, multiplying the above equations together member by member, we have

$$(30) \quad f(q) = [\text{II}] \frac{1}{1 - q^{2h}},$$

or

$$(31) \quad A = \frac{1}{(1 - q^2)(1 - q^4)(1 - q^6) \dots}.$$

Substituting this value in [equation \(8\)](#), Chap. VIII, we have, after making  $u = 0$ ,

$$\begin{aligned} (1 - q)^2(1 - q^3)^2(1 - q^5)^2 \dots &= \frac{1 - 2q + 2q^4 - 2q^9 + \dots}{(1 - q^2)(1 - q^4)(1 - q^6) \dots} \\ &= \frac{\Theta(0)}{(1 - q^2)(1 - q^4)(1 - q^6) \dots}. \end{aligned}$$

(See [equation \(9\)](#), Chap. VIII.)

Transposing one of the series of products from the left-hand member, we get

$$(1 - q)(1 - q^3) \dots = \frac{\Theta(0)}{(1 - q)(1 - q^2)(1 - q^3)(1 - q^4) \dots}.$$

Introducing on both sides of the equation the factors  $1 - q^2$ ,  $1 - q^4$ ,  $1 - q^6$ , etc., we get

$$\begin{aligned} &(1 - q)(1 - q^2)(1 - q^3)(1 - q^4) \dots \\ &= \Theta(0) \frac{1 - q^2}{1 - q} \cdot \frac{1 - q^4}{1 - q^2} \cdot \frac{1 - q^6}{1 - q^3} \cdot \frac{1 - q^8}{1 - q^4} \dots \\ &= \Theta(0)(1 + q)(1 + q^2)(1 + q^3) \dots; \end{aligned}$$

whence

$$(32) \quad \Theta(0) = \frac{(1-q)(1-q^2)(1-q^3)\cdots}{(1+q)(1+q^2)(1+q^3)\cdots}.$$

Resuming [equation \(20\)](#), Chap. VII, and dividing both members of the equation by  $u$ , we have

$$\frac{\operatorname{sn} u}{u} = \frac{2\sqrt[4]{q}}{\sqrt{k}} \frac{\sin \frac{\pi u}{2K}}{u} [\Pi] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

This, for  $u = 0$ , since the limiting value of  $\frac{\operatorname{sn} u}{u}$  for  $u = 0$  is 1, and of  $\frac{\sin \frac{\pi u}{2K}}{u}$  for  $x = 0$  is  $\frac{\pi}{2K'}$ , becomes

$$1 = \frac{\sqrt[4]{q}}{\sqrt{k}} \cdot \frac{\pi}{K} \cdot \frac{(1-q^2)^2(1-q^4)^2(1-q^6)^2\cdots}{(1-q)^2(1-q^3)^2(1-q^5)^2\cdots},$$

or

$$(33) \quad \frac{\sqrt{k}K}{\pi \sqrt[4]{q}} = \left[ \frac{(1-q^2)(1-q^4)(1-q^6)\cdots}{(1-q)(1-q^3)(1-q^5)\cdots} \right]^2.$$

Further, from [equation \(21\)](#), Chap. VII, for  $u = 0$ , we have

$$(34) \quad \frac{\sqrt{k}}{2\sqrt{k'}\sqrt[4]{q}} = \left[ \frac{(1+q^2)(1+q^4)(1+q^6)\cdots}{(1-q)(1-q^3)(1-q^5)\cdots} \right]^2.$$

The quotient of these two equations gives

$$(35) \quad \frac{2\sqrt{k'}K}{\pi} = \left[ \frac{(1-q^2)(1-q^4)(1-q^6)\cdots}{(1+q^2)(1+q^4)(1+q^6)\cdots} \right]^2;$$

or, substituting the value of  $\sqrt{k'}$  from eqs. (18) and (19), Chap. VII,

$$(36) \quad \frac{2k'K}{\pi} = \left[ \frac{(1-q)(1-q^2)(1-q^3)\cdots}{(1+q)(1+q^2)(1+q^3)\cdots} \right]^2.$$

Comparing this with equation (32), we easily get

$$(37) \quad \Theta(0) = \sqrt{\frac{2k'K}{\pi}}.$$

From equation (9), Chap. VIII, making  $u = K$ , we get

$$(38) \quad \Theta(K) = 1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \cdots.$$

Making  $z = 0$  in equation (24), Chap. IX, we have

$$(39) \quad \Theta_1(0) = 1 + 2q + 2q^4 + 2q^9 + \cdots.$$

This might also have been derived from eq. (38) by observing that

$$\Theta(0 + K) = \Theta_1(0) = \Theta(K).$$

Knowing  $\Theta(0)$ , it is easy to deduce  $\Theta(K)$  and  $H(K)$ .

From equation (7) we have

$$\operatorname{dn} u = \sqrt{k'} \frac{\Theta(u + K)}{\Theta(u)}.$$

Making  $u = 0$ , we have, since  $\operatorname{dn}(0) = 1$ ,

$$(40) \quad \Theta(K) = \frac{\Theta(0)}{\sqrt{k'}}.$$

From equation (5) we get, in the same manner,

$$(41) \quad H(K) = \sqrt{\frac{k'}{k}} \Theta(0).$$

From eq. (12), Chap. IX, we have

$$(41)^* \quad \operatorname{dn} u = \sqrt{1 - k^2 \sin^2 \phi} = \sqrt{k'} \frac{\Theta_1(u)}{\Theta(u)};$$

and putting  $x = \frac{\pi u}{2K}$ , we have

$$(42) \quad \frac{\operatorname{dn} u}{\sqrt{k'}} = \frac{1 + 2q \cos 2x + 2q^4 \cos 4x + 2q^9 \cos 6x + \dots}{1 - 2q \cos 2x + 2q^4 \cos 4x - 2q^9 \cos 6x + \dots}.$$

Putting

$$(42)^* \quad \frac{\operatorname{dn} u}{\sqrt{k'}} = \cot \gamma,$$

we have

$$\frac{\cot \gamma - 1}{\cot \gamma + 1} = \tan(45^\circ - \gamma) = 2q \frac{\cos 2x + q^8(4 \cos^3 2x - 3 \cos 2x) + \dots}{1 + q^4(4 \cos^2 2x - 2)};$$

whence

$$(43) \quad \cos 2x = \frac{\tan(45^\circ - \gamma)[1 + q^4(4 \cos^2 2x - 2)]}{2q} - q^8(4 \cos^3 2x - 3 \cos 2x) - \dots,$$

and approximately,

$$(44) \quad \cos 2x = \frac{\tan(45^\circ - \gamma)}{2q}.$$

From equations (37) and (40), Chap. IX, we have

$$(45) \quad x = \frac{u}{\Theta^2(K)};$$

whence

$$(46) \quad u = x\Theta^2(K).$$

# CHAPTER X.

## ELLIPTIC INTEGRALS OF THE SECOND ORDER.

FROM Chap. I, [equation \(19\)](#), we have

$$E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi = \int_0^\phi \Delta\phi \cdot d\phi.$$

From this we have

$$E(\phi) + E(\psi) = \int_0^\phi \Delta\phi \cdot d\phi + \int_0^\psi \Delta\phi \cdot d\phi.$$

Put

$$(1) \quad E\phi + E\psi = S.$$

Differentiating, we get

$$(2) \quad \Delta\phi \cdot d\phi + \Delta\psi \cdot d\psi = dS.$$

But we have, Chap. II, [equation \(2\)](#),

$$\frac{d\phi}{\Delta\phi} + \frac{d\psi}{\Delta\psi} = 0,$$

or

$$(3) \quad \Delta\psi \cdot d\phi + \Delta\phi \cdot d\psi = 0.$$

Adding equations [\(2\)](#) and [\(3\)](#), we get

$$(4) \quad (\Delta\phi + \Delta\psi)(d\phi + d\psi) = dS.$$

Substituting  $\cos \mu$  from eq. (5), in eq. (5)\*, Chap. II, we get

$$(5) \quad \begin{cases} \Delta\phi = \frac{\sin \phi \cos \psi \Delta\mu + \cos \phi \sin \psi}{\sin \mu}, \\ \Delta\psi = \frac{\sin \psi \cos \phi \Delta\mu + \cos \psi \sin \phi}{\sin \mu}; \end{cases}$$

whence

$$(6) \quad \Delta\phi \pm \Delta\psi = \frac{\Delta\mu \pm 1}{\sin \mu} \sin(\phi \pm \psi).$$

Substituting in equation (4), we have

$$(7) \quad \begin{aligned} dS &= \frac{\Delta\mu + 1}{\sin \mu} \sin(\phi + \psi) d(\phi + \psi) \\ &= -\frac{\Delta\mu + 1}{\sin \mu} d \cos(\phi + \psi). \end{aligned}$$

Integrating equation (7), we have

$$E\phi + E\psi = \frac{\Delta\mu + 1}{\sin \mu} [C - \cos(\phi + \psi)].$$

The constant of integration,  $C$ , is determined by making  $\phi = 0$ ; in this case  $\psi = \mu$ ,  $E\phi = 0$ ,  $E\psi = E\mu$ , and  $S = E\mu$ ; whence

$$E\mu = \frac{\Delta\mu + 1}{\sin \mu} (C - \cos \mu),$$

and by subtraction,

$$E\phi + E\psi - E\mu = \frac{\Delta\mu + 1}{\sin \mu} (\cos \mu - \cos \phi \cos \psi + \sin \phi \sin \psi).$$

But, Chap. II, eq. (5),

$$\cos \mu - \cos \phi \cos \psi = -\sin \phi \sin \psi \Delta \mu;$$

whence

$$E\phi + E\psi - E\mu = \frac{1 - \Delta^2 \mu}{\sin \mu} \sin \phi \sin \psi$$

whence

$$(8) \quad E\phi + E\psi = E\mu + k^2 \sin \phi \sin \psi \sin \mu.$$

When  $\phi = \psi$ , we have

$$(9) \quad E\mu = 2E\phi - k^2 \sin^2 \phi \sin \mu.$$

But in that case

$$(10) \quad \cos \mu = \cos^2 \phi - \sin^2 \phi \Delta \mu;$$

whence

$$(11) \quad \sin \phi = \sqrt{\frac{1 - \cos \mu}{1 + \Delta \mu}}.$$

Let  $\phi, \phi_{\frac{1}{2}}, \phi_{\frac{1}{4}}$ , etc., be such values as will satisfy the equations

$$(12) \quad \begin{aligned} E\phi &= 2E\phi_{\frac{1}{2}} - k^2 \sin^2 \phi_{\frac{1}{2}} \sin \phi, \\ E\phi_{\frac{1}{2}} &= 2E\phi_{\frac{1}{4}} - k^2 \sin^2 \phi_{\frac{1}{4}} \sin \phi_{\frac{1}{2}}, \\ &\text{etc.} \qquad \qquad \text{etc.} \end{aligned}$$

Assume an auxiliary angle  $\gamma$ , such that

$$(13) \quad \sin \gamma = k \sin \phi;$$





since  $\sin \phi = 1 - \frac{\phi^3}{3!} + \frac{\phi^5}{5!} - \text{etc.}$ ,

$$\begin{aligned} \Delta\phi &= (1 - k^2 \sin^2 \phi)^{\frac{1}{2}} \\ &= 1 - \frac{k^2}{2} \left( \phi - \frac{\phi^3}{6} \right)^2 - \frac{k^4}{8} \left( \phi - \frac{\phi^3}{6} \right)^4 + \dots \\ &= 1 - \frac{k^2}{2} \phi^2 + \left( \frac{k^2}{6} - \frac{k^4}{8} \right) \phi^4. \end{aligned}$$

Whence

$$\begin{aligned} (17) \quad Ek\phi_{\frac{1}{2^n}} &= \int_0^{\phi_n} \Delta\phi_{\frac{1}{2^n}} d\phi \\ &= \phi_{\frac{1}{2^n}} - \frac{k^2}{6} \phi_{\frac{1}{2^n}}^3 + \frac{k^2(4-3k^2)}{120} \phi_{\frac{1}{2^n}}^5. \end{aligned}$$

Substituting in [eq. \(16\)](#) the numerical values derived from [equations \(15\)](#) and [\(17\)](#), we are enabled to determine the value of  $E\phi$ .

Landen's Transformation can also be applied to Elliptic Integrals of this class.

From [eq. \(11\)](#), Chap. IV, we get, by easy transformation,

$$(18) \quad \sin^2 2\phi = \sin^2 \phi_1 (1 + k_0 + 2k_0 \cos 2\phi).$$

From this we easily get

$$\begin{aligned} 2k_0 \cos 2\phi \sin^2 \phi_1 &= \sin^2 2\phi - \sin^2 \phi_1 - k_0^2 \sin^2 \phi_1 \\ &= 1 - \cos^2 2\phi - \sin^2 \phi_1 - k_0^2 \sin^2 \phi_1 \\ &= \Delta^2 k_0 \phi_1 - \sin^2 \phi_1 - \cos^2 2\phi; \end{aligned}$$

whence

$$\cos^2 2\phi + 2k_0 \sin^2 \phi_1 \cos 2\phi = \Delta^2 k_0 \phi_1 - \sin^2 \phi_1;$$

and from this,

$$\begin{aligned}
 \cos 2\phi &= -k_0 \sin^2 \phi_1 \pm \sqrt{\Delta^2 k_0 \phi_1 - \sin^2 \phi_1 + k_0^2 \sin^4 \phi_1} \\
 (19) \qquad &= \cos \phi_1 \Delta k_0 \phi_1 - k_0 \sin^2 \phi_1;
 \end{aligned}$$

whence, also,

$$\begin{aligned}
 1 - \cos^2 2\phi &= 1 - \cos^2 \phi_1 \Delta^2 \phi_1 + 2k \sin^2 \phi_1 \cos \phi_1 \Delta k_0 \phi_1 - k_0^2 \sin^4 \phi_1 \\
 &= \sin^2 \phi_1 (1 + k_0^2 \cos^2 \phi_1 + 2k_0 \cos \phi_1 \Delta k_0 \phi_1 - k_0^2 \sin^2 \phi_1)
 \end{aligned}$$

and

$$(20) \qquad \sin 2\phi = \sin \phi_1 (\Delta k_0 \phi_1 + k_0 \cos \phi_1).$$

Differentiating [equation \(19\)](#), we get

$$2 \sin 2\phi \frac{d\phi}{d\phi_1} = \sin \phi_1 \frac{(k_0 \cos \phi_1 + \Delta k_0 \phi_1)^2}{\Delta k_0 \phi_1}.$$

Dividing this by [equation \(20\)](#), we have

$$\frac{2d\phi}{d\phi_1} = \frac{k_0 \cos \phi_1 + \Delta k_0 \phi_1}{\Delta k_0 \phi_1}.$$

But from [\(19\)](#), and [eq. \(6\)](#), Chap. IV,

$$\begin{aligned}
 k^2 \sin^2 \phi &= \frac{k^2(1 - \cos 2\phi)}{2} \\
 &= \frac{2k_0}{(1 + k_0)^2} \{1 + k_0 \sin^2 \phi_1 - \cos \phi_1 \Delta k_0 \phi_1\};
 \end{aligned}$$

whence

$$\Delta k \phi = \frac{\Delta k_0 \phi_1 + k_0 \cos \phi_1}{1 + k_0},$$

and

$$2\Delta k \phi \cdot \frac{d\phi}{d\phi_1} = \frac{(k_0 \cos \phi_1 + \Delta k_0 \phi_1)^2}{(1 + k_0)\Delta k_0 \phi_1},$$

and

$$d\phi \Delta k \phi = \frac{d\phi_1}{\Delta k_0 \phi_1} \cdot \frac{(k_0 \cos \phi_1 + \Delta k_0 \phi_1)^2}{2(1 + k_0)}.$$

This gives immediately, by integration,

$$\begin{aligned} Ek\phi &= \frac{1}{2(1 + k_0)} \int \frac{d\phi_1}{\Delta k_0 \phi_1} \{k_0 \cos \phi_1 + \Delta k_0 \phi_1\}^2 \\ &= \frac{1}{2(1 + k_0)} \int \frac{d\phi_1}{\Delta k_0 \phi_1} \{2\Delta^2 k_0 \phi_1 + 2k_0 \cos \phi_1 \Delta k_0 \phi_1 - k_1'^2\} \\ (21) \quad &= \frac{Ek_0 \phi_1}{1 + k_0} + \frac{k_0 \sin \phi_1}{1 + k_0} - \frac{1}{2}(1 - k_0)Fk_0 \phi_1. \end{aligned}$$

Thus the value of  $Ek\phi$  is made to depend upon  $Ek_0\phi_1$  (containing a smaller modulus and a larger amplitude), and upon the integral of the first class,  $Fk_0\phi_1$ ;  $k_0$ ,  $\phi_1$ , etc., being determined by the formulæ (6) to (12) of Chap. IV.

By successive applications of equation (21),  $Ek\phi$  may be made to depend ultimately upon  $Ek_{0n}\phi_n$ , where  $k_{0n}$  approximates to zero and  $Ek_{0n}\phi_n$  to  $\phi_n$ .

Or, by reversing, it may be made to depend upon  $Ek_n\phi_{0n}$ , where  $k_n$  approximates to unity and  $Ek_n\phi_{0n}$  to  $-\cos \phi_{0n}$ .

To facilitate this, assume

$$Gk\phi = Ek\phi - Fk\phi.$$

Subtracting from equation (21) the equation

$$Fk\phi = \frac{1 + k_0}{2} Fk_0 \phi_1 \text{ (see eq. (13), Chap. IV),}$$

we have

$$Gk\phi = \frac{1}{1+k_0}(Gk_0\phi_1 + k_0 \sin \phi_1 - k_0 Fk_0 \phi_1).$$

Repeated applications of this give

$$Gk_0\phi_1 = \frac{1}{1+k_{00}}(Gk_{00}\phi_2 + k_{00} \sin \phi_2 - k_{00} Fk_{00} \phi_2),$$

.....

$$Gk_{0(n-1)}\phi_{n-1} = \frac{1}{1+k_{0n}}(Gk_{0n}\phi_n + k_{0n} \sin \phi_n - k_{0n} Fk_{0n} \phi_n).$$

Whence

$$(22) \quad Gk\phi = \sum_n \left\{ \frac{k_{0n}(\sin \phi_n - Fk_{0n}\phi_n)}{\left[\prod_n^1\right](1+k_{0n})} \right\} + \frac{Gk_{0n}\phi_n}{\left[\prod_n^1\right](1+k_{0n})}.$$

But since (compare [eq. \(13\)](#), Chap. IV)

$$Fk\phi = \frac{Fk_{0n}\phi_n \left[\prod_n^1\right](1+k_{0n})}{2^n},$$

or

$$(23) \quad \frac{Fk_{0n}\phi_n}{\left[\prod_n^1\right](1+k_{0n})} = \frac{2^n Fk\phi}{\left[\prod_n^1\right](1+k_{0n})^2};$$

and since, also, (compare [eq. \(6\)](#), Chap. IV,)

$$\frac{k_{0(n-1)}^2}{k_{0n}} = \frac{2^2}{(1+k_{0n})^2},$$

we have

$$\begin{aligned}
 (24) \quad \frac{2^n k_{0n}}{\left[\prod_n^1\right] (1 + k_{0n})^2} &= \frac{k_{0n}}{2^n} \left[\prod_n^1\right] \frac{k_{0(n-1)}^2}{k_{0n}} \\
 &= \frac{k_{0n}}{2^n} \left[\prod_n^1\right] \frac{k_{0(n-1)}}{k_{0n}} \left[\prod_n^1\right] k_{0(n-1)} \\
 &= \frac{k_{0n}}{2^n} \cdot \frac{k}{k_0} \cdot \frac{k_0}{k_{00}} \cdots \frac{k_{0(n-1)}}{k_{0n}} \cdot k \left[\prod_n^2\right] k_{0(n-1)} \\
 &= \frac{k^2}{2^n} \left[\prod_n^2\right] k_{0(n-1)}.
 \end{aligned}$$

Substituting these values in [equation \(22\)](#), and neglecting the term containing  $Gk_{0n}\phi_n$  since, carried to its limiting value,

$$\begin{aligned}
 Gk_{0n}\phi_n &= Ek_{0n}\phi_n - Fk_{0n}\phi_n \\
 &= \phi_n - \phi_n = 0, \quad (n = \text{limiting value,})
 \end{aligned}$$

we have

$$\begin{aligned}
 (25) \quad Gk\phi &= \sum_n \left\{ \frac{k\sqrt{k_{0n}} \sin \phi_n \left[\prod_n^2\right] \sqrt{k_{0(n-1)}} - k^2 \left[\prod_n^2\right] k_{0(n-1)}}{2^n} \right\} \\
 &= k \left[ \frac{\sqrt{k_0}}{2} \sin \phi_1 + \frac{\sqrt{k_0 k_{00}}}{2^2} \sin \phi_2 + \frac{\sqrt{k_0 k_{00} k_{03}}}{2^3} \sin \phi_3 + \cdots \right] \\
 &\quad - \frac{k^2}{2} \left[ 1 + \frac{k_0}{2} + \frac{k_0 k_{00}}{2^2} + \frac{k_0 k_{00} k_{03}}{2^3} + \cdots \right];
 \end{aligned}$$

whence

$$(26) \quad Ek\phi = Fk\phi \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_0}{2} + \frac{k_0k_{00}}{2^2} + \dots \right) \right] \\ + k \left[ \frac{\sqrt{k_0}}{2} \sin \phi_1 + \frac{\sqrt{k_0k_{00}}}{2^2} \sin \phi_2 + \frac{\sqrt{k_0k_{00}k_{03}}}{2^3} \sin \phi_3 + \dots \right].$$

From eq. (3), Chap. V, we see that when  $\phi = \frac{\pi}{2}$ ,

$$\phi_n = 2^{n-1}\pi.$$

Substituting these values in equation (26), we have for a complete Elliptic Integral of the second class,

$$(27) \quad E\left(k, \frac{\pi}{2}\right) = \\ F\left(k, \frac{\pi}{2}\right) \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_0}{2} + \frac{k_0k_{00}}{2^2} + \frac{k_0k_{00}k_{03}}{2^3} + \dots \right) \right].$$

In a similar manner we could have found the formula for  $E(k, \phi)$  in terms of an increasing modulus, viz.,

$$(28) \quad E(k, \phi) = F(k, \phi) \left[ 1 + k \left( 1 + \frac{2}{k_1} + \frac{2^2}{k_1k_2} + \frac{2^3}{k_1k_2k_3} + \dots \right. \right. \\ \left. \left. + \frac{2^{n-2}}{k_1k_2 \dots k_{n-2}} - \frac{2^{n-1}}{k_1k_2 \dots k_{n-1}} \right) \right] \\ - k \left[ \sin \phi + \frac{2}{\sqrt{k}} \sin \phi_1 + \frac{2^2}{\sqrt{kk_1}} \sin \phi_2 + \dots \right. \\ \left. + \frac{2^{n-1}}{\sqrt{kk_1 \dots k_{n-2}}} \sin \phi_{n-1} - \frac{2^n}{\sqrt{kk_1 \dots k_{n-1}}} \sin \phi_n \right].$$

# CHAPTER XI.

## ELLIPTIC INTEGRALS OF THE THIRD ORDER.

THE Elliptic Integral of the third order is

$$(1) \quad \Pi(n, k, \phi) = \int_0^\phi \frac{d\phi}{(1 + n \sin^2 \phi) \Delta\phi}.$$

Put

$$(2) \quad \Pi(\phi) + \Pi(\psi) = S;$$

whence we have immediately

$$(3) \quad dS = \frac{d\phi}{(1 + n \sin^2 \phi) \Delta\phi} + \frac{d\psi}{(1 + n \sin^2 \psi) \Delta\psi}.$$

But, [eq. \(2\)](#), Chap. II,

$$(4) \quad \frac{d\phi}{\Delta\phi} + \frac{d\psi}{\Delta\psi} = 0;$$

whence

$$(5) \quad \begin{aligned} dS &= \left( \frac{1}{1 + n \sin^2 \phi} - \frac{1}{1 + n \sin^2 \psi} \right) \frac{d\phi}{\Delta\phi} \\ &= \frac{n(\sin^2 \psi - \sin^2 \phi)}{(1 + n \sin^2 \phi)(1 + n \sin^2 \psi)} \cdot \frac{d\phi}{\Delta\phi}. \end{aligned}$$

From [equation \(8\)](#), Chap. X, we get by differentiation, since  $\sigma$  (or  $\mu$ ) is constant,

$$\Delta\phi \cdot d\phi + \Delta\psi \cdot d\psi = k^2 \sin \sigma d(\sin \phi \sin \psi),$$

or, from [equation \(3\)](#),

$$(\sin^2 \psi - \sin^2 \phi) \frac{d\phi}{\Delta\phi} = \sin \sigma d(\sin \phi \sin \psi).$$

This, introduced into [equation \(5\)](#), gives

$$dS = \frac{n \sin \sigma d(\sin \phi \sin \psi)}{1 + n(\sin^2 \phi + \sin^2 \psi) + n^2 \sin^2 \phi \sin^2 \psi}.$$

Put

$$\sin \phi \sin \psi = q, \quad \sin^2 \phi + \sin^2 \psi = p;$$

whence

$$(6) \quad dS = \frac{n \sin \sigma dq}{1 + np + n^2 p^2}.$$

From [equation \(5\)](#), Chap. II, we have

$$\cos \sigma = \cos \phi \cos \psi - \sin \phi \sin \psi \Delta\sigma,$$

from which we easily get

$$\begin{aligned} (\cos \sigma + q\Delta\sigma)^2 &= \cos^2 \phi \cos^2 \psi \\ &= (1 - \sin^2 \phi)(1 - \sin^2 \psi) \\ &= 1 - p + q^2, \end{aligned}$$

and thence

$$\begin{aligned} p &= 1 + q^2 - (\cos \sigma + q \Delta\sigma)^2 \\ &= \sin^2 \sigma - 2 \cos \sigma \Delta\sigma q + k^2 \sin^2 \sigma \cdot q^2. \end{aligned}$$



This, substituted in eq. (6), gives

$$\begin{aligned} dS &= \frac{n \sin \sigma dq}{1 + n \sin^2 \sigma - 2n \cos \sigma \Delta \sigma q + n(n + k^2 \sin^2 \sigma)q^2} \\ &= \frac{n \sin \sigma dq}{A - 2Bq + Cq^2}, \end{aligned}$$

where

$$\begin{aligned} A &= 1 + n \sin^2 \sigma, \\ B &= n \cos \sigma \Delta \sigma, \\ C &= nk^2 \sin^2 \sigma + n^2. \end{aligned}$$

From this we get

$$S = n \sin \sigma \int \frac{dq}{A - 2Bq + Cq^2} + \text{Const.}$$

In order to determine the constant of integration we must observe that for  $\phi = 0$ ,  $\psi = \sigma$  and  $q = 0$ ; whence

$$\Pi\sigma = n \sin \sigma \int_{q=0} \frac{dq}{A - 2Bq + Cq^2} + \text{Const.};$$

whence

$$S = \Pi\sigma + n \sin \sigma \int_0^q \frac{dq}{A - 2Bq + Cq^2},$$

or

$$(7) \quad \Pi\phi + \Pi\psi = \Pi\sigma + n \sin \sigma \int_0^q \frac{dq}{A - 2Bq + Cq^2}.$$

But we have

$$\begin{aligned}
 dS &= \frac{CM dq}{AC - B^2 + (Cq - B)^2} \\
 &= \frac{CM}{AC - B^2} \cdot \frac{dq}{1 + \left( \frac{Cq - B}{\sqrt{AC - B^2}} \right)^2} \\
 &= \frac{M}{\sqrt{AC - B^2}} \cdot \frac{\frac{C dq}{\sqrt{AC - B^2}}}{1 + \left( \frac{Cq - B}{\sqrt{AC - B^2}} \right)^2}
 \end{aligned}$$

where  $M = n \sin \sigma$ .

The integral of the second member is

$$\frac{M}{\sqrt{AC - B^2}} \tan^{-1} \frac{Cq - B}{\sqrt{AC - B^2}};$$

whence

$$\int_0^q dS = S_1 = \frac{M}{\sqrt{AC - B^2}} \left[ \tan^{-1} \frac{Cq - B}{\sqrt{AC - B^2}} + \tan^{-1} \frac{B}{\sqrt{AC - B^2}} \right];$$

or, since

$$\begin{aligned}
 \tan^{-1} x + \tan^{-1} y &= \tan^{-1} \frac{x + y}{1 - xy}, \\
 S_1 &= \frac{M}{\sqrt{AC - B^2}} \tan^{-1} \frac{q\sqrt{AC - B^2}}{A - Bq}.
 \end{aligned}$$

Substituting the values of  $A$ ,  $B$ ,  $C$  and  $M$ , we have

$$\begin{aligned}
 AC - B^2 &= n(1 + n - \Delta^2 \sigma)(1 + n \sin^2 \sigma) - n^2 \cos^2 \sigma \Delta^2 \sigma \\
 &= n(1 + n - \Delta^2 \sigma + n(1 + n) \sin^2 \sigma - n \Delta^2 \sigma) \\
 &= n(1 + n)(1 - \Delta^2 \sigma + n \sin^2 \sigma) \\
 &= n(1 + n)(k^2 + n) \sin^2 \sigma;
 \end{aligned}$$

and putting

$$\frac{(1+n)(k^2+n)}{n} = \Omega,$$

we have

$$\sqrt{AC - B^2} = n\sqrt{\Omega} \sin \sigma.$$

Substituting these values in [eq. \(7\)](#), we have

$$\begin{aligned} & \Pi(n, k, \phi) + \Pi(n, k, \psi) - \Pi(n, k, \sigma) = S_1 \\ & = \frac{1}{\sqrt{\Omega}} \tan^{-1} \frac{n\sqrt{\Omega} \sin \phi \sin \psi \sin \sigma}{1 + n \sin^2 \sigma - n \sin \phi \sin \psi \cos \sigma \Delta \sigma}. \end{aligned}$$

# CHAPTER XII.

## NUMERICAL CALCULATIONS. $q$ .

### CALCULATION OF THE VALUE OF $q$ .

FROM [eq. \(7\)](#), Chap. IX, we have

$$\operatorname{dn} u = \sqrt{k'} \frac{\Theta(u + K)}{\Theta(u)};$$

whence, [eq. \(9\)](#), Chap. IV, [eqs. \(27\)](#) and [\(39\)](#), Chap. IX,

$$(1) \quad \begin{aligned} \sqrt{\cos \theta} &= \frac{1 - 2q + 2q^4 - 2q^9 + 2q^{16} - \dots}{1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \dots} \\ &= 1 - 4q + 8q^2 - 16q^3 + 32q^4 - 56q^5 + \dots \end{aligned}$$

The first five terms of this series can be represented by

$$\sqrt{\cos \theta} = \frac{1 - 2q}{1 + 2q}.$$

From this we get

$$(2) \quad q = \frac{1}{2} \cdot \frac{1 - \sqrt{\cos \theta}}{1 + \sqrt{\cos \theta}},$$

which is exact up to the term containing  $q^5$ .

Or we can deduce a more exact formula as follows: From [eq. \(1\)](#),

$$\begin{aligned} \frac{1 + \sqrt{\cos \theta}}{1 - \sqrt{\cos \theta}} &= \frac{\sqrt{1 + \tan^2 \frac{1}{2}\theta} + \sqrt{1 - \tan^2 \frac{1}{2}\theta}}{\sqrt{1 + \tan^2 \frac{1}{2}\theta} - \sqrt{1 - \tan^2 \frac{1}{2}\theta}} \\ &= \frac{1 + 2q^4 + 2q^{16} + \dots}{2q + 2q^9 + 2q^{25} + \dots}; \end{aligned}$$



When  $\theta$  approaches  $90^\circ$ ,  $\tan \frac{\theta}{2}$  differs little from unity, and the series in eq. (5) is not very converging, but  $q$  can be calculated by means of eq. (6), Chap. VII, viz.,

$$q = e^{-\frac{\pi K'}{K}}, \quad q' = e^{-\frac{\pi K}{K'}}.$$

By comparing these equations with eqs. (6) and (9), Chap. IV, we see that if

$$q = f(k) = f(\theta),$$

then

$$q' = f(k') = f(90^\circ - \theta).$$

Therefore, having  $\theta$ , we can from its complement,  $90^\circ - \theta$ , find  $q'$  by eq. (5), and thence  $q$  by the following process. We have

$$\frac{1}{q} = e^{\frac{\pi K'}{K}}, \quad \frac{1}{q'} = e^{\frac{\pi K}{K'}};$$

whence

$$\begin{aligned} \log \frac{1}{q} \log \frac{1}{q'} &= M^2 \pi^2 = 1.8615228, \\ (6) \quad \log \log \frac{1}{q} + \log \log \frac{1}{q'} &= 0.2698684, \end{aligned}$$

by which we can deduce  $q$  from  $q'$ .

EXAMPLE. Let  $\theta = 79^\circ 36' 14''$ . To find  $q$ .

$$90^\circ - \theta = 10^\circ 23' 46''.$$

By eq. (5) we get

$$\log q' = 7.3156316, \quad \log \frac{1}{q'} = 2.6843684,$$

$$\text{and } \log \log \frac{1}{q'} = .4288421;$$

and by eq. (6),

$$\log \log \frac{1}{q} = 9.8410263;$$

whence

$$\log q = 1.3065321.$$

When  $k' = k = \cos 45^\circ = \frac{1}{2}\sqrt{2}$ , eq. (6) becomes

$$(7) \quad \log \frac{1}{q} = M\pi = 1.3643763; \quad (k = k';)$$

whence

$$\begin{aligned} \log q &= 2.6356237, \\ q &= 0.0432138. \end{aligned} \quad (k = k'.)$$

EXAMPLE. Given  $\theta = 10^\circ 23' 46''$ . Find  $q$ .

*Ans.*  $\log q = 7.3156316$ .

EXAMPLE. Given  $\theta = 82^\circ 45'$ . Find  $q$ .

*Ans.*  $\log q = 9.37919$ .

# CHAPTER XIII.

## NUMERICAL CALCULATIONS. K.

### CALCULATION OF THE VALUE OF K.

WE have already found from eq. (37), Chap. IX,

$$(1) \quad \Theta(0) = \sqrt{\frac{2k'K}{\pi}},$$

and from eq. (40), same chapter,

$$(2) \quad \Theta(K) = \frac{\Theta(0)}{\sqrt{k'}} = \sqrt{\frac{2K}{\pi}}.$$

But, eqs. (38) and (27), Chap. IX,

$$\begin{aligned} \Theta(K) &= 1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \dots, \\ \Theta(0) &= 1 - 2q + 2q^4 - 2q^9 + 2q^{16} - \dots; \end{aligned}$$

whence, eq. (2),

$$(3) \quad K = \frac{\pi}{2}(1 + 2q + 2q^4 + 2q^9 + \dots)^2.$$

By adding eqs. (1) and (2) we get

$$\Theta(0) + \Theta(K) = \sqrt{\frac{2K}{\pi}}(1 + \sqrt{k'});$$



whence

$$\begin{aligned}
 K &= \frac{\pi}{2} \left( \frac{\Theta(0) + \Theta(K)}{1 + \sqrt{k'}} \right)^2 \\
 &= \frac{\pi}{2} \left[ \frac{2(1 + 2q^4 + 2q^{16} + \dots)}{1 + \sqrt{k'}} \right]^2 \\
 (4) \quad &= \frac{\pi}{2} \left( \frac{2}{1 + \sqrt{k'}} \right)^2 (1 + 2q^4 + 2q^{16} + \dots)^2.
 \end{aligned}$$

EXAMPLE. Let  $k = \sin \theta = \sin 19^\circ 30'$ . Required  $K$ .

*First Method.* By eq. (3).

By eq. (5), Chap. XII, we find  $\log q = 8.6356236$ . Applying eq. (3), using only two terms of the series, we have

$$\begin{aligned}
 1 + 2q &= 1.0147662 \\
 \log(1 + 2q) &= 0.0063660 \\
 2\log(1 + 2q) &= 0.0127320 \\
 \log \frac{\pi}{2} &= 0.1961199 \\
 \hline
 \log K &= 0.2088519 \\
 K &= 1.615101
 \end{aligned}$$

*Second Method.* By eq. (4).

Equation (4) may be written, neglecting  $q^4$ ,

$$K = \frac{\pi}{2} \left( \frac{1 + \sqrt{\cos \theta}}{2} \right)^{-2};$$

whence

$$\begin{aligned} \log \cos \theta &= 9.9743466, \\ \log \sqrt{\cos \theta} &= 9.9871733, \\ 1 + \sqrt{\cos \theta} &= 1.9708973, \\ \frac{1 + \sqrt{\cos \theta}}{2} &= 0.98544865; \end{aligned}$$

and

$$\begin{aligned} \log K &= 0.2088519, \\ K &= 1.615101, \end{aligned}$$

the same result as above.

*Third Method.* By eq. (7), Chap. V.

$\begin{aligned} \theta &= 19^\circ 30' \\ \frac{1}{2}\theta &= 9^\circ 45' \\ \log \tan \frac{1}{2}\theta &= 9.235103 \\ \log \cos \frac{1}{2}\theta &= 9.993681 \\ \left. \begin{aligned} \log \tan^2 \frac{1}{2}\theta \\ \log \sin \theta \end{aligned} \right\} &= 8.470206 \\ \theta_0 &= 1^\circ 41' 31''.1 \end{aligned}$		$\begin{aligned} \theta_0 &= 1^\circ 41' 31''.1 \\ \frac{1}{2}\theta_0 &= 0^\circ 50' 45''.5 \\ \log \cos \frac{1}{2}\theta_0 &= 9.999953 \end{aligned}$
---	--	--

$$\log \cos^2 \frac{1}{2}\theta = 9.987362$$

$$\log \cos^2 \frac{1}{2}\theta_0 = 9.999906$$

$$\underline{9.987268}$$

$$\log \frac{\pi}{2} = 0.196120$$

$$\log K = 0.208852$$

$\theta_{00}$  is not calculated, as it is evident that its cosine will be 1.

EXAMPLE. Given  $k = \sin 75^\circ$ . Find  $K$ .

By eq. (7), Chap. V.

From eqs. (14<sub>1</sub>), Chap. IV, we find

$$k = \sin \theta = \sin 75^\circ \qquad \log = 9.9849438$$

$$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta = \tan^2 37^\circ 30' \\ \sin \theta_0 = \sin 36^\circ 4' 16'' .47 \end{array} \right\} \qquad 9.7699610$$

$$k_{00} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_0 = \tan^2 18^\circ 2' 8'' .235 \\ \sin \theta_{00} = \sin 6^\circ 5' 9'' .38 \end{array} \right\} \qquad 9.0253880$$

$$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_{00} = \tan^2 3^\circ 2' 34'' .69 \\ \sin \theta_{03} = \sin 9' 42'' .90 \end{array} \right\} \qquad 7.4511672$$

	log	2 log	a. c. 2 log
$\cos \frac{1}{2}\theta = \cos 37^\circ 30'$	9.8994667	9.7989334	0.2010666
$\cos \frac{1}{2}\theta_0 = \cos 18^\circ 2'.13725$	9.9781184	9.9562368	0.0437632
$\cos \frac{1}{2}\theta_{02} = \cos 3^\circ 2'.57817$	9.9993873	9.9987746	0.0012254
$\cos \frac{1}{2}\theta_{03} = \cos 4'.8575$	9.9999995	9.9999990	0.0000010
			<hr style="width: 100%; border: 0.5px solid black;"/> 0.2460562
$\frac{\pi}{2}$		$\frac{\pi}{2}$	<hr style="width: 100%; border: 0.5px solid black;"/> .1961199
		log K =	0.4421761
		$K = 2.768064$	<i>Ans.</i>

EXAMPLE. Given  $k = \sin 45^\circ$ . Find  $K$ .  
 Method of [eq. \(7\)](#), Chap. V.  
 From [eqs. \(14<sub>1</sub>\)](#), Chap. IV, we have

	log
$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta = \tan^2 22^\circ 30' \\ \sin \theta_0 = \sin 9^\circ 52'.75683 \end{array} \right\}$	9.2344486
$k_{00} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_0 = \tan^2 4^\circ 56'.37841 \\ \sin \theta_{00} = \sin 25'.679 \end{array} \right\}$	7.8733009
$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_{00} = \tan^2 12'.3395 \\ \sin \theta_{03} = \sin 0'.05 \end{array} \right\}$	5.1445523

$$\text{a. c. } \log \cos^2 \frac{1}{2}\theta \qquad 0.0687694$$

$$\text{a. c. } \log \cos^2 \frac{1}{2}\theta_0 \qquad 0.0032320$$

$$\text{a. c. } \log \cos^2 \frac{1}{2}\theta_{00} \qquad 0.0000060$$

$$\log \frac{\pi}{2} \qquad \underline{0.1961199}$$

$$\log K = 0.2681273$$

$$K = 1.8540747 \quad \text{Ans.}$$

EXAMPLE. Given  $\theta = 63^\circ 30'$ . Find  $K$ .

$$\text{Ans. } \log K = 0.3539686.$$

EXAMPLE. Given  $\theta = 34^\circ 30'$ . Find  $K$ .

$$\text{Ans. } K = 1.72627.$$

# CHAPTER XIV.

## NUMERICAL CALCULATIONS. $u$

### CALCULATION OF THE VALUE OF $u$ .

WHEN  $\theta^\circ = \sin^{-1} k < 45^\circ$ .

EXAMPLE. Let  $\phi = 30^\circ$ ,  $k = \sin 45^\circ$ . Find  $u$ .

*First Method.* Eq. (23), Chap. IV, and eqs. (14<sub>1</sub>), (14<sub>2</sub>), (14<sub>3</sub>), Chap. IV.  
By equations (14<sub>1</sub>),

$$\frac{\theta}{2} = 22^\circ 30';$$

$$\log \tan \frac{\theta}{2} = 9.6172243;$$

$$\log \tan^2 \frac{\theta}{2} = 9.2344486 = \log k_0 = \log \sin \theta_0;$$

$$\theta_0 = 9^\circ 52' 45''.41;$$

$$\log \tan \frac{\theta_0}{2} = 8.9366506;$$

$$\log \tan^2 \frac{\theta_0}{2} = 7.8733012 = \log k_{00} = \log \sin \theta_{00};$$

$$\theta_{00} = 0^\circ 25' 40''.7;$$

$$\log \tan^2 \frac{\theta_{00}}{2} = 5.144552 = \log k_{03}.$$

By equations (14<sub>2</sub>),

$$\phi = 30^\circ$$

$$\log \tan \phi = 9.761439$$

$$\log \cos \theta = 9.849485$$

---


$$\log \tan(\phi_1 - \phi) = 9.610924$$

$$\phi_1 - \phi = 22^\circ 12' 27''.56$$

$$\phi_1 = 52^\circ 12' 27''.56$$

$$\log \tan \phi_1 = 0.110438$$

$$\log \cos \theta_0 = 9.993512$$

---


$$\log \tan(\phi_2 - \phi_1) = 0.103949$$

$$\phi_2 - \phi_1 = 51^\circ 47' 32''.59$$

$$\phi_2 = 104^\circ 0' 0''.15$$

$$\log \tan \phi_2 = 0.603228$$

$$\log \cos \theta_{00} = 9.999988$$

---


$$\log \tan(\phi_3 - \phi_2) = 0.603216$$

$$\phi_3 - \phi_2 = 104^\circ 0' 1''.5$$

$$\phi_3 = 208^\circ 0' 1''.65$$

Since  $\frac{\phi_2}{4} = 26^\circ 0' 0''.04$  and  $\frac{\phi_3}{8} = 26^\circ 0' 0''.21$ , we need not calcu-

late  $\phi_4$ .

$$\frac{\phi_3}{8} = 93600'' .21.$$

Reducing this to radians, we have

$$\log \frac{\phi_3}{8} = 9.656852.$$

Substituting in eq. (23), Chap. IV, we have, since  $\cos \theta_{03} = 1$ ,

$$\text{a. c. } \log \cos \theta = 0.150515$$

$$\log \cos \theta_0 = 9.993512$$

$$\log \cos \theta_{00} = 9.999988$$

---


$$0.144014$$

$$0.072007 = \log \sqrt{\frac{\cos \theta_0 \cos \theta_{00}}{\cos \theta}}$$

$$\log \frac{\phi_3}{8} = 9.656852$$

$$\log u = 9.728859$$

$$u = 0.535623, \quad \text{Ans.}$$

When  $\theta = \sin^{-1} k > 45^\circ$ .

EXAMPLE. Given  $k = \sin 75^\circ$ ,  $\tan \phi = \sqrt{\frac{2}{\sqrt{3}}}$ . To find  $F(k, \phi)$ .

*First Method. Bisected Amplitudes.*



By [equations \(24\)](#) and [\(25\)](#), Chap. IV, we get

$$\begin{array}{ll}
 \phi & = 47^\circ 3' 30''.91, \\
 \phi_{\frac{1}{2}} & = 25^\circ 36' 5''.64, & \beta & = 45^\circ; \\
 \phi_{\frac{1}{4}} & = 13^\circ 6' 30''.98, & \beta_0 & = 24^\circ 40' 10''.94; \\
 \phi_{\frac{1}{8}} & = 6^\circ 35' 40''.74, & \beta_{00} & = 12^\circ 39' 15''.83; \\
 \phi_{\frac{1}{16}} & = 3^\circ 18' 8''.75, & \beta_{03} & = 6^\circ 22' 8''.40; \\
 \phi_{\frac{1}{32}} & = 1^\circ 39' 7''.43, & \beta_{04} & =
 \end{array}$$

Substituting in [equation \(26\)](#), Chap. IV, we have

$$\begin{aligned}
 F(k, \phi) &= 32 \times 1^\circ 39' 7''.43 \\
 &= 52^\circ 51' 58''.03 \\
 &= 0.9226878. \quad \text{Ans.}
 \end{aligned}$$

*Second Method.* [Equation \(29\)](#), Chap. IV.

From [equations \(18<sub>3</sub>\)](#), Chap. IV, we have

	log
$k = \cos \eta = \cos 15^\circ 0' 0''.00$	9.9849438
$k' = \sin \eta = \sin 15^\circ 0' 0''.00$	9.4129962
$k'_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\eta = \tan^2 7^\circ 30' 0''.00 \\ \sin \eta_0 = \sin 0^\circ 59' 35''.25 \end{array} \right\}$	8.2388582
$k_1 = \cos \eta_0 = \cos 0^\circ 59' 35''.25$	9.9999348
$k'_{00} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\eta_0 = \tan^2 0^\circ 29' 47''.62 \\ \sin \eta_{00} = \sin 0^\circ 0' 15''.49 \end{array} \right\}$	5.8757219
$k_2 = \cos \eta_{00} = \cos 0^\circ 0' 15''.49$	0.0000000
$k'_{03} = \left(\frac{1}{2}k'_{00}\right)^2$	1.1493838

From [equations \(18<sub>2</sub>\)](#), Chap. IV, we get

$$\begin{aligned} \phi &= 47^\circ 3' 30''.95; \\ 2\phi_0 - \phi &= 45^\circ; \\ \phi_0 &= 46^\circ 1' 45''.475; \\ \phi_{02} &= 46^\circ 1' 29''.41; \\ \phi_{03} &= 46^\circ 1' 29''.41; \\ 45^\circ + \frac{1}{2}\phi_3 &= 68^\circ 0' 44''.705. \end{aligned}$$

Substituting these values in [eq. \(29\)](#), Chap. IV, we get

$$\begin{aligned} F(k, \phi) &= \sqrt{\frac{k_1}{k}} \cdot \frac{1}{M} \cdot \log \tan 68^\circ 0' 44''.705 \\ &= 0.9226877. \quad \text{Ans.} \end{aligned}$$

Third Method. Equation (23), Chap. IV.

From equations (14<sub>1</sub>), Chap. IV, we have

		log
$k = \sin \theta$	$= \sin 75^\circ 0' 0''$	9.9849438
$k' = \cos \theta$	$= \cos 75^\circ$	9.4129962
$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta \\ \sin \theta_0 \end{array} \right.$	$= \left\{ \begin{array}{l} \tan^2 37^\circ 30' \\ \sin 36^\circ 4' 16''.47 \end{array} \right.$	9.7699610
$k'_1 = \cos \theta_0$		9.9075648
$k_{02} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_0 \\ \sin \theta_{00} \end{array} \right.$	$= \left\{ \begin{array}{l} \tan^2 18^\circ 2' 8''.235 \\ \sin 6^\circ 5' 9''.38 \end{array} \right.$	9.0253880
$k'_2 = \cos \theta_{00}$		9.9975452
$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_{00} \\ \sin \theta_{03} \end{array} \right.$	$= \left\{ \begin{array}{l} \tan^2 3^\circ 2' 34''.69 \\ \sin 9' 42''.90 \end{array} \right.$	7.4511672
$k'_3 = \cos \theta_{03}$		9.9999982
$k_{04} = \left( \frac{1}{2} k_{03} \right)^2$		4.3002761
$k'_4 =$		0.0000000

From equations (14<sub>2</sub>), Chap. IV, we have

$$\begin{aligned} \phi &= 47^\circ 3' 30''.94; \\ \phi_1 &= 62^\circ 36' 3''.10; \\ \phi_2 &= 119^\circ 55' 47''.67; \\ \phi_3 &= 240^\circ 0' 0''.19; \\ \phi_4 &= 480^\circ 0' 0''. \end{aligned}$$

Therefore the limit of  $\phi$ ,  $\frac{\phi_1}{2}$ ,  $\frac{\phi_2}{4}$ , or  $\frac{\phi_n}{2^n}$  is  $30^\circ = \frac{\pi}{6}$ .

Substituting these values in eq. (23), Chap. IV, we have

$$\begin{aligned} F(k, \phi) &= \sqrt{\frac{k'_1 k'_2 k'_3 k'_4}{k'}} \cdot \frac{\pi}{6} \\ &= 0.9226874. \quad \text{Ans.} \end{aligned}$$

EXAMPLE. Given  $\phi = 30^\circ$ ,  $k = \sin 89^\circ$ . Find  $u$ .

Method of eq. (28), Chap. IV.

From eqs. (18<sub>1</sub>) we find

$$k_1 = \sin \theta_1 \quad \text{and} \quad \tan^2 \frac{1}{2} \theta_1 = k = \sin \theta,$$

from which we find that  $k_1 = 1$  as far as seven decimal places.

From eqs. (18<sub>2</sub>) we have

$$\begin{aligned} \sin \phi &= 9.6989700 \\ k &= \underline{9.9999338} \\ \sin(2\phi_0 - \phi) &= 9.6989038 \\ 2\phi_0 - \phi &= 29^\circ 59'.69733 \\ 2\phi_0 &= 59^\circ 59'.69733 \\ 45^\circ + \frac{1}{2}\phi_0^* &= 59^\circ 59'.92433 \\ \log \left( 45^\circ + \frac{1}{2}\phi_0 \right) &= 0.2385385 \end{aligned}$$

From eqs. (18<sub>3</sub>), Chap. IV, we have

$$k = \cos \eta = \cos 1^\circ, \quad \frac{1}{2}\eta = 30'.$$

---

\*Since  $k_1 = 1$ ,  $\phi_{00} = \phi_0$ , and we need not carry the calculation further.

Substituting in eq. (28), Chap. IV, we have

$$\text{a. c. } \log \cos \frac{1}{2}\eta \quad 0.0000330$$

$$\log \log \left( 45^\circ + \frac{1}{2}\phi_0 \right) \quad 9.3775585$$

$$\text{a. c. } \log M \quad \underline{0.3622157}$$

$$\log F(k, \phi) = 9.7398072$$

$$F(k, \phi) = 0.549297. \quad \text{Ans.}$$

EXAMPLE. Given  $\phi = 79^\circ$ ,  $k = 0.25882$ . Find  $u$ .

$$\text{Ans. } u = 0.39947.$$

EXAMPLE. Given  $\phi = 37^\circ$ ,  $k = 0.86603$ . Find  $u$ .

$$\text{Ans. } u = 0.68141.$$

# CHAPTER XV.

## NUMERICAL CALCULATIONS. $\phi$ .

EXAMPLE. Given  $u = 1.368407$ ,  $\theta = 38^\circ$ . Find  $\phi$ .

*First Method.* From eqs. (46) and (41)\*, Chap. IX, we have

$$u = x\Theta^2(K),$$
$$\Delta\phi = \sqrt{k'} \frac{\Theta_1(x)}{\Theta(x)}.$$

From equations (5), Chap. XII, and (38), Chap. IX, we have

$$\log q = 8.4734187$$

$$\log \Theta^2(K) = 0.0501955$$

$$\log u = \underline{0.1362153}$$

$$\log x = 0.0860198$$

$$x = 69^\circ 50' 46''.12$$

From equations (23) and (24), Chap. IX, we get

$$\log \Theta_1(x) = 9.9798368$$

$$\log \Theta(x) = \underline{0.0192687}$$

$$9.9605681$$

$$\log \sqrt{k'} = \underline{9.9482661}$$

$$\log \Delta\phi = 9.9088342 = \log \sin \lambda$$

But

$$\begin{aligned} k^2 \sin^2 \phi &= 1 - \Delta^2 \phi, \\ k \sin \phi &= \cos \lambda; \end{aligned}$$

whence

$$\begin{aligned} \log \cos \lambda &= 9.7675483 \\ \log k &= \underline{9.7893420} \\ \log \sin \phi &= 9.9782063 \\ \phi &= 72^\circ. \quad \text{Ans.} \end{aligned}$$

*Second Method.* From eq. (1), Chap. VI.

From eqs. (14<sub>1</sub>) Chap. IV, we find

		log
$k_0 =$	$\left\{ \begin{array}{ll} \tan^2 \frac{1}{2}\theta & = \tan^2 19^\circ \\ \sin \theta_0 & = \sin 6^\circ 48' .54569 \end{array} \right\}$	9.0739438
		$\cos \theta_0$
$k_{00} =$	$\left\{ \begin{array}{ll} \tan^2 \frac{1}{2}\theta_0 & = \tan^2 3^\circ 24' .2784 \\ \sin \theta_{00} & = \sin 12' .16659 \end{array} \right\}$	7.5488952
		$\cos \theta_{00}$
$k_{03} =$	$\left\{ \begin{array}{ll} \tan^2 \frac{1}{2}\theta_{00} & = \tan^2 6' .08329 \\ \sin \theta_{03} & \end{array} \right\}$	4.4957316
		$\cos \theta_{03}$

Substituting these values in eq. (1), Chap. VI, we have

$\log \cos \theta_0$	9.9969260
$\log \cos \theta_{00}$	9.9999974
	9.9969234
$\log \sqrt{\cos \theta_0 \cos \theta_{00}}$	9.9984617
a. c. log “ “	0.0015383
$\log u$	.1362153
$\log \sqrt{\cos \theta}$	9.9482660
$\log 2^3$	.9030900*
a. c. log $\sqrt{\cos \theta_0 \cos \theta_{00}}$	0.0015383
	0.9891096
	2.2418773
$\log \phi_3^*$	2.7472323
$\phi_3$	558° 46'.140

---

\* $n$  is taken equal to 3, because  $\cos \theta_{03} = 1$ .



Whence, by [equations \(1\)](#) of Chap. VI, we get

$$\begin{aligned} k_{03} \log &= 4.4957316 \\ \sin \phi_3 & \frac{9.5075232_n}{4.0032548_n} \\ \sin(2\phi_2 - \phi_3) & \\ 2\phi_2 - \phi_3 &= -0'.00346 \\ \phi_2 &= 279^\circ 23'.06827 \end{aligned}$$

$$\begin{aligned} k_{00} \log &= 7.5488952 \\ \sin \phi_2 & \frac{9.9941484_n}{7.5430436_n} \\ \sin(2\phi_1 - \phi_2) & \\ 2\phi_1 - \phi_2 &= -12'.0039 \\ \phi_1 &= 139^\circ 35'.5321 \end{aligned}$$

$$\begin{aligned} k_0 \log &= 9.0739438 \\ \sin \phi_1 & \frac{9.8117249}{8.8856687} \\ \sin(2\phi - \phi_1) & \\ 2\phi - \phi_1 &= 4^\circ 24'.467 \\ \phi &= 71^\circ 59'.9999 \\ &= 72^\circ. \quad \text{Ans.} \end{aligned}$$

EXAMPLE. Given  $u = 2.41569$ ,  $\theta = 80^\circ$ . Find  $\phi$ .

Ans.  $\phi = 82^\circ$ .

EXAMPLE. Given  $u = 1.62530$ ,  $k = \frac{1}{2}$ . Find  $\phi$ .

Ans.  $\phi = 87^\circ$ .

# CHAPTER XVI.

## NUMERICAL CALCULATIONS. $E(k, \phi)$ .

*First Method.* By Chap. X, eqs. (15), (16), and (17).

EXAMPLE. Given  $k = 0.9327$ ,  $\phi = 80^\circ$ . Find  $E(k, \phi)$ .

By eq. (15), Chap. X,

$$\phi = 80^\circ;$$

$$\gamma = 67^\circ 44'. ;$$

$$\phi_{\frac{1}{2}} = 50^\circ 43'.6,$$

$$\gamma_{\frac{1}{2}} = 46^\circ 40'.4;$$

$$\phi_{\frac{1}{4}} = 27^\circ 48'.5,$$

$$\gamma_{\frac{1}{4}} = 26^\circ 0'.1;$$

$$\phi_{\frac{1}{8}} = 14^\circ 16'.7,$$

$$\gamma_{\frac{1}{8}} = 13^\circ 24'.0;$$

$$\phi_{\frac{1}{16}} = 7^\circ 11'.3,$$

$$\gamma_{\frac{1}{16}} = 6^\circ 45'.2;$$

$$\phi_{\frac{1}{32}} = 3^\circ 36'.0,$$

$$\log \sin \gamma_{\frac{1}{32}} = 8.77094;$$

$$\phi_{\frac{1}{32}} = 0.062831.$$

$$\therefore \phi_{\frac{1}{32}}^5 < 0.0000001.$$

Whence, by eq. (17),

$$\begin{aligned}
 E(k, \phi_{\frac{1}{32}}) &= 0.062794 \\
 \sin \phi \sin^2 \gamma_{\frac{1}{2}} &= 0.52116 \\
 2 \sin \phi_{\frac{1}{2}} \sin^2 \gamma_{\frac{1}{4}} &= 0.29757 \\
 4 \sin \phi_{\frac{1}{4}} \sin^2 \gamma_{\frac{1}{8}} &= 0.10023 \\
 8 \sin \phi_{\frac{1}{8}} \sin^2 \gamma_{\frac{1}{16}} &= 0.02728 \\
 16 \sin \phi_{\frac{1}{16}} \sin^2 \gamma_{\frac{1}{32}} &= 0.00697 \\
 &\quad \underline{\hspace{1.5cm}} \\
 &\quad 0.95321
 \end{aligned}$$

Hence, by eq. (16),

$$\begin{aligned}
 E(k, \phi) &= 32E(k, \phi_{\frac{1}{32}}) - 0.95321 \\
 &= 2.0094 - 0.9532 = 1.0562.
 \end{aligned}$$

*Second Method.* By Chap. X, eq. (26).

EXAMPLE. Given  $k = \sin 75^\circ$ ,  $\tan \phi = \sqrt{\frac{2}{\sqrt{3}}}$ . Find  $E(k, \phi)$ .

From eqs. (14<sub>1</sub>), Chap. IV, we have

$k = \sin \theta = \sin 75^\circ 0' 0''$	log = 9.9849438
$k' = \cos \theta = \cos 75^\circ$	9.4129962
$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta = \tan^2 37^\circ 30' \\ \sin \theta_0 = \sin 36^\circ 4' 16''.47 \end{array} \right\}$	9.7699610
$k'_1 = \cos \theta_0$	9.9075648
$k_{02} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_0 = \tan^2 18^\circ 2' 8''.235 \\ \sin \theta_{00} = \sin 6^\circ 5' 9''.38 \end{array} \right\}$	9.0253880
$k'_2 = \cos \theta_{00}$	9.9975452
$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_{00} = \tan^2 3^\circ 2' 34''.69 \\ \sin \theta_{03} = \sin 9' 42''.90 \end{array} \right\}$	7.4511672
$k'_3 = \cos \theta_{03}$	9.9999982
$k_{04} = \left( \frac{1}{2} k_{03} \right)^2$	4.3002761
$k'_4 =$	0.0000000

From eqs. (14<sub>2</sub>), Chap. IV, we have

$$\begin{aligned} \phi &= 47^\circ 3' 30''.94; \\ \phi_1 &= 62^\circ 36' 3''.10; \\ \phi_2 &= 119^\circ 55' 47''.67; \\ \phi_3 &= 240^\circ 0' 0''.19. \end{aligned}$$

Applying eq. (26), Chap. X, we have

$$\begin{array}{r} k^2 \log = 9.9698876 \\ \text{a. c. 2} \quad \underline{9.6989700} \\ \quad \quad \quad 9.6688576 \quad .4665064 \end{array}$$

$k_0$	9.7699610	
a. c. 2	<u>9.6989700</u>	
	9.1377886	.1373373
$k_{00}$	9.0253880	
a. c. 2	<u>9.6989700</u>	
	7.8621466	.0072802
$k_{03}$	7.4511672	
a. c. 2	<u>9.6989700</u>	
	5.0132838	<u>.0000103</u>
		.6111342

$$1 - .6111342 = 0.3888658.$$

From eq. (23), Chap. IV, we find  $F(k, \phi) = 0.9226874$ .

Hence

$$\begin{aligned}
 F(k, \phi) \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_0}{2} + \dots \right) \right] &= 0.3588016 \\
 \frac{k\sqrt{k_0}}{2} \sin \phi_1 &= 0.3290186 \\
 \frac{k\sqrt{k_0 k_{00}}}{4} \sin \phi_2 &= 0.0522872 \\
 \frac{k\sqrt{k_0 k_{02} k_{03}}}{8} \sin \phi_3 &= -0.0013888 \\
 \frac{k\sqrt{k_0 \dots k_{04}}}{16} \sin \phi_4 &= \underline{0.0000010} \\
 &= 0.3799180
 \end{aligned}$$

Whence

$$E(k, \phi) = 0.3588016 + 0.3799180 = 0.7387196. \quad \text{Ans.}$$

EXAMPLE. Given  $k = \sin 75^\circ$ . Find  $E\left(k, \frac{\pi}{2}\right)$ .

From Example 2, Chap. XIII, we find

$$\log F\left(k, \frac{\pi}{2}\right) = 0.4421761$$

$$\log 0.3888658 = 1.5897998$$

$$\log E\left(k, \frac{\pi}{2}\right) = \overline{0.0319759}$$

$$E\left(k, \frac{\pi}{2}\right) = 1.076405. \quad \text{Ans.}$$

EXAMPLE. Given  $k = \sin 30^\circ$ ,  $\phi = 81^\circ$ . Find  $E(k, \phi)$ .

$$\text{Ans. } E(k, \phi) = 1.33124.$$

EXAMPLE. Find  $E(\sin 80^\circ, 55^\circ)$ .

$$\text{Ans. } 0.82417.$$

EXAMPLE. Find  $E\left(\sin 27^\circ, \frac{\pi}{2}\right)$ .

$$\text{Ans. } 1.48642.$$

EXAMPLE. Find  $E(\sin 19^\circ, 27^\circ)$ .

$$\text{Ans. } 0.46946.$$

# CHAPTER XVII.

## APPLICATIONS.

### RECTIFICATION OF THE LEMNISCATE.

THE polar equation of the Lemniscate is  $r = a\sqrt{\cos 2\theta}$ , referred to the centre as the origin. From this we get

$$\frac{dr}{d\theta} = -\frac{a \sin 2\theta}{\sqrt{\cos 2\theta}};$$

whence the length of the arc measured from the vertex to any point whose co-ordinates are  $r$  and  $\theta$

$$\begin{aligned} s &= \int \left\{ \left( \frac{dr}{d\theta} \right)^2 + r^2 \right\}^{\frac{1}{2}} d\theta = a \int \left\{ \frac{\sin^2 2\theta}{\cos 2\theta} + \cos 2\theta \right\}^{\frac{1}{2}} d\theta \\ &= a \int \frac{d\theta}{\sqrt{\cos 2\theta}} = a \int \frac{d\theta}{\sqrt{1 - 2\sin^2 \theta}}. \end{aligned}$$

Let  $\cos 2\theta = \cos^2 \phi$ , whence

$$\begin{aligned} s &= a \int \frac{\frac{d\theta}{d\phi} d\phi}{\cos \phi} = a \int \frac{\sin \phi d\phi}{\sqrt{1 - \cos^4 \phi}} \\ &= a \int_0^\phi \frac{d\phi}{\sqrt{1 + \cos^2 \phi}} = \frac{a}{\sqrt{2}} \int_0^\phi \frac{d\phi}{\sqrt{1 - \frac{1}{2} \sin^2 \phi}} \\ &= \frac{a}{\sqrt{2}} F \left( \frac{1}{\sqrt{2}}, \phi \right). \end{aligned}$$

Since  $r = a\sqrt{\cos 2\theta} = a \cos \phi$ , the angle  $\phi$  can be easily constructed by describing upon the axis  $a$  of the Lemniscate a semicircle, and then

revolving the radius vector until it cuts this semicircle. In the right-angled triangle of which this is one side, and the axis the hypotenuse,  $\phi$  is evidently the angle between the axis and the revolved position of the radius vector.

## RECTIFICATION OF THE ELLIPSE.

Since the equation of the ellipse is  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , we can assume  $x = a \sin \phi$ ,  $y = b \cos \phi$ , so that  $\phi$  is the complement of the *eccentric angle*. Hence

$$\begin{aligned} s &= \int \sqrt{dx^2 + dy^2} = a \int d\phi \sqrt{1 - e^2 \sin^2 \phi} \\ &= aE(e, \phi), \end{aligned}$$

in which  $e$ , the eccentricity of the ellipse, is the modulus of the Elliptic Integral.

The length of the Elliptic Quadrant is

$$s' = aE\left(e, \frac{\pi}{2}\right).$$

EXAMPLE. The equation of an ellipse is

$$\frac{x^2}{16.81} + \frac{y^2}{16} = 1;$$

required the length of an arc whose abscissas are 1.061162 and 4.100000: of the quadrantal arc.

*Ans.* 5.18912; 6.36189.

## RECTIFICATION OF THE HYPERBOLA.

On the curve of the hyperbola, construct a straight line perpendicular to the axis  $x$ , and at a distance from the centre equal to the projection



of  $b$ , the transverse axis, upon the asymptote, i.e. equal to  $\frac{b^2}{\sqrt{a^2 + b^2}}$ . Join the projection of the given point of the hyperbola on this line with the centre. The angle which this joining line makes with the axis of  $x$  we will call  $\phi$ . If  $y$  is the ordinate of the point on the hyperbola, then evidently

$$y = \frac{b^2 \tan \phi}{\sqrt{a^2 + b^2}},$$

and

$$x = \frac{a}{\cos \phi} \sqrt{1 - \frac{a^2 \sin^2 \phi}{a^2 + b^2}} = \frac{a}{\cos \phi} \sqrt{1 - \frac{1}{e^2} \sin^2 \phi};$$

whence

$$\begin{aligned} s &= \int \sqrt{dx^2 + dy^2} = \frac{b^2}{c} \int_0^\phi \frac{d\phi}{\cos^2 \phi \sqrt{1 - \frac{1}{e^2} \sin^2 \phi}} \\ &= \frac{b^2}{c} \int_0^\phi \frac{d\phi}{\cos^2 \phi \sqrt{1 - k^2 \sin^2 \phi}}. \end{aligned}$$

But

$$\begin{aligned} d(\tan \phi \sqrt{1 - k^2 \sin^2 \phi}) &= d\phi \sqrt{1 - k^2 \sin^2 \phi} + d\phi \frac{1 - k^2}{\sqrt{1 - e^2 \sin^2 \phi}} \\ &\quad - \frac{1 - k^2}{\cos^2 \phi \sqrt{1 - e^2 \sin^2 \phi}} d\phi. \end{aligned}$$

Consequently

$$\begin{aligned} s &= \frac{b^2}{c} \int_0^\phi \frac{d\phi}{\cos^2 \phi \sqrt{1 - k^2 \sin^2 \phi}} \\ &= \frac{b^2}{c} F(k, \phi) - cE(k, \phi) + c \tan \phi \Delta(k, \phi) \\ &= \frac{b^2}{ae} F\left(\frac{1}{e}, \phi\right) - aeE\left(\frac{1}{e}, \phi\right) + ae \tan \phi \Delta\left(\frac{1}{e}, \phi\right). \end{aligned}$$

EXAMPLE. Find the length of the arc of the hyperbola

$$\frac{x^2}{20.25} - \frac{y^2}{400} = 1$$

from the vertex to the point whose ordinate is  $\frac{40}{2.05} \tan 15^\circ$ .

*Ans.* 5.231184.

EXAMPLE. Find the length of the arc of the hyperbola

$$\frac{x^2}{144} - \frac{y^2}{81} = 100$$

from the vertex to the point whose ordinate is 0.6.

*Ans.* 0.6582.

End of the Project Gutenberg EBook of Elliptic Functions, by Arthur L. Baker

\*\*\* END OF THIS PROJECT GUTENBERG EBOOK ELLIPTIC FUNCTIONS \*\*\*

\*\*\*\*\* This file should be named 31076-pdf.pdf or 31076-pdf.zip \*\*\*\*\*

This and all associated files of various formats will be found in:

<http://www.gutenberg.org/3/1/0/7/31076/>

Produced by Andrew D. Hwang, Brenda Lewis and the Online Distributed Proofreading Team at <http://www.pgdp.net> (This file was produced from images from the Cornell University Library: Historical Mathematics Monographs collection.)

Updated editions will replace the previous one--the old editions will be renamed.

Creating the works from public domain print editions means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg-tm electronic works to protect the PROJECT GUTENBERG-tm concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for the eBooks, unless you receive specific permission. If you do not charge anything for copies of this eBook, complying with the rules is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. They may be modified and printed and given away--you may do practically ANYTHING with public domain eBooks. Redistribution is subject to the trademark license, especially commercial redistribution.

\*\*\* START: FULL LICENSE \*\*\*

THE FULL PROJECT GUTENBERG LICENSE  
PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg-tm mission of promoting the free

distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase "Project Gutenberg"), you agree to comply with all the terms of the Full Project Gutenberg-tm License (available with this file or online at <http://gutenberg.org/license>).

## Section 1. General Terms of Use and Redistributing Project Gutenberg-tm electronic works

1.A. By reading or using any part of this Project Gutenberg-tm electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg-tm electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg-tm electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. "Project Gutenberg" is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg-tm electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg-tm electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg-tm electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation ("the Foundation" or PGLAF), owns a compilation copyright in the collection of Project Gutenberg-tm electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is in the public domain in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg-tm mission of promoting free access to electronic works by freely sharing Project Gutenberg-tm works in compliance with the terms of

this agreement for keeping the Project Gutenberg-tm name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg-tm License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg-tm work. The Foundation makes no representations concerning the copyright status of any work in any country outside the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg-tm License must appear prominently whenever any copy of a Project Gutenberg-tm work (any work on which the phrase "Project Gutenberg" appears, or with which the phrase "Project Gutenberg" is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at [www.gutenberg.org](http://www.gutenberg.org)

1.E.2. If an individual Project Gutenberg-tm electronic work is derived from the public domain (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase "Project Gutenberg" associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg-tm trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg-tm electronic work is posted with the permission of the copyright holder, your use and distribution

must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg-tm License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg-tm License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg-tm.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg-tm License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg-tm work in a format other than "Plain Vanilla ASCII" or other format used in the official version posted on the official Project Gutenberg-tm web site ([www.gutenberg.org](http://www.gutenberg.org)), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original "Plain Vanilla ASCII" or other form. Any alternate format must include the full Project Gutenberg-tm License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg-tm works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg-tm electronic works provided that

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg-tm works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg-tm trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you

prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."

- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg-tm License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg-tm works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg-tm works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg-tm electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from both the Project Gutenberg Literary Archive Foundation and Michael Hart, the owner of the Project Gutenberg-tm trademark. Contact the Foundation as set forth in Section 3 below.

#### 1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread public domain works in creating the Project Gutenberg-tm collection. Despite these efforts, Project Gutenberg-tm electronic works, and the medium on which they may be stored, may contain "Defects," such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the "Right of Replacement or Refund" described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg-tm trademark, and any other party distributing a Project Gutenberg-tm electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH F3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS' WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone



providing copies of Project Gutenberg-tm electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg-tm electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg-tm work, (b) alteration, modification, or additions or deletions to any Project Gutenberg-tm work, and (c) any Defect you cause.

## Section 2. Information about the Mission of Project Gutenberg-tm

Project Gutenberg-tm is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need, are critical to reaching Project Gutenberg-tm's goals and ensuring that the Project Gutenberg-tm collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg-tm and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation web page at <http://www.pglaf.org>.

## Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Its 501(c)(3) letter is posted at <http://pglaf.org/fundraising>. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's principal office is located at 4557 Melan Dr. S.

Fairbanks, AK, 99712., but its volunteers and employees are scattered throughout numerous locations. Its business office is located at 809 North 1500 West, Salt Lake City, UT 84116, (801) 596-1887, email [business@pglaf.org](mailto:business@pglaf.org). Email contact links and up to date contact information can be found at the Foundation's web site and official page at <http://pglaf.org>

For additional contact information:

Dr. Gregory B. Newby  
Chief Executive and Director  
[gnewby@pglaf.org](mailto:gnewby@pglaf.org)

#### Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg-tm depends upon and cannot survive without wide spread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine readable form accessible by the widest array of equipment including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit <http://pglaf.org>

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg Web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: <http://pglaf.org/donate>

Section 5. General Information About Project Gutenberg-tm electronic works.

Professor Michael S. Hart is the originator of the Project Gutenberg-tm concept of a library of electronic works that could be freely shared with anyone. For thirty years, he produced and distributed Project Gutenberg-tm eBooks with only a loose network of volunteer support.

Project Gutenberg-tm eBooks are often created from several printed editions, all of which are confirmed as Public Domain in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our Web site which has the main PG search facility:

<http://www.gutenberg.org>

This Web site includes information about Project Gutenberg-tm, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.